

SHARP DEGREE BOUNDS FOR SUM-OF-SQUARES CERTIFICATES ON PROJECTIVE CURVES

GRIGORIY BLEKHERMAN, GREGORY G. SMITH, AND MAURICIO VELASCO

ABSTRACT. Given a real projective curve with homogeneous coordinate ring R and a nonnegative homogeneous element $f \in R$, we bound the degree of a nonzero homogeneous sum-of-squares $g \in R$ such that the product fg is again a sum of squares. Better yet, our degree bounds only depend on geometric invariants of the curve and we show that there exists smooth curves and nonnegative elements for which our bounds are sharp. We deduce the existence of a multiplier g from a new Bertini Theorem in convex algebraic geometry and prove sharpness by deforming rational Harnack curves on toric surfaces. Our techniques also yield similar bounds for multipliers on surfaces of minimal degree, generalizing Hilbert's work on ternary forms.

1. OVERVIEW OF RESULTS

Certifying that a polynomial is nonnegative remains a central problem in real algebraic geometry and optimization. The quintessential certificate arises from multiplying a given polynomial by a second polynomial, that is already known to be positive, and expressing the product as a sum of squares. Although the Positivstellensatz guarantees that suitable multipliers exist over any semi-algebraic set, tight bounds on the degree of multipliers are exceptionally rare. Our primary aim is to produce sharp degree bounds for sum-of-squares multipliers on real projective curves. In reaching this goal, the degree bounds also reveal a surprising consonance between real and complex algebraic geometry.

To be more explicit, fix an embedded real projective curve $X \subset \mathbb{P}^n$ that is nondegenerate and totally real—not contained in a hyperplane and with Zariski-dense real points. Let R be its \mathbb{Z} -graded coordinate ring and let $r(X)$ denote the least integer i such that the Hilbert polynomial and function of X agree at all integers greater than or equal to i . For $j \in \mathbb{N}$, we write $P_{X,2j} \subset R_{2j}$ and $\Sigma_{X,2j} \subset R_{2j}$ for the cone of nonnegative elements in R_{2j} and the cone of sums of squares of elements from R_j , respectively. Our first result gives a sharp degree bound on sum-of-squares multipliers in terms of the fundamental geometric invariants of X .

Theorem 1.1. *If $X \subset \mathbb{P}^n$ is a nondegenerate totally-real projective curve of degree d and arithmetic genus p_a and $f \in P_{X,2j}$ is a nonnegative element of positive degree then, for all $k \in \mathbb{N}$ with $k \geq \max\{r(X), \frac{2p_a}{d}\}$, there is a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$. Conversely, for all $n \geq 2$ and for all $j \geq 2$, there exist totally-real smooth curves $X \subset \mathbb{P}^n$ and nonnegative elements $f \in P_{X,2j}$ such that, for all $k < \max\{r(X), \frac{2p_a}{d}\}$ and for all nonzero $g \in \Sigma_{X,2k}$, we have $fg \notin \Sigma_{X,2j+2k}$.*

Remarkably, the uniform degree bound on the multiplier g is determined by the complex geometry of the curve X . It is independent of both the degree of the nonnegative element f and the Euclidean topology of the real points in X .

Date: 26 May 2016.

2010 Mathematics Subject Classification. 14P05; 12D15, 14H45.

Our approach also applies to higher-dimensional varieties that are arithmetically Cohen–Macaulay, but it is most effective on some surfaces. A subvariety $X \subset \mathbb{P}^n$ has minimal degree if it is non-degenerate and $\deg(X) = 1 + \text{codim}(X)$. Theorem 1.1 in [BSV] establishes that $P_{X,2} = \Sigma_{X,2}$ if and only if X is a totally-real variety of minimal degree. Evocatively, this equivalence leads to a characterization of the varieties over which multipliers of degree zero suffice. Building on this framework and generalizing Hilbert’s work [Hil] on ternary forms, our second result gives degree bounds for sum-of-squares multipliers on surfaces of minimal degree.

Theorem 1.2. *If $X \subset \mathbb{P}^n$ is a totally-real surface of minimal degree and $f \in P_{X,2j}$ is a nonnegative element of positive degree, then there is a nonzero $g \in \Sigma_{j^2-j}$ such that $fg \in \Sigma_{X,j^2+j}$. Conversely, if $X \subset \mathbb{P}^n$ is a totally-real surface of minimal degree and $j \geq 2$, then there exist nonnegative elements $f \in P_{X,2j}$ such that, for all $k < j - 2$ and for all nonzero $g \in \Sigma_{X,2k}$, we have $fg \notin \Sigma_{X,2j+2k}$.*

Unlike curves, Theorem 1.2 shows that the minimum degree of a sum-of-squares multiplier g depends intrinsically on the degree of the nonnegative element f . The sharpness of the upper or lower bounds on these surfaces is an intriguing open problem.

Motivated by its relation to Hilbert’s Seventeenth Problem, we obtain slightly better degree bounds when the totally-real surface is \mathbb{P}^2 ; see Example 4.18 and Example 5.17 for the details. Specifically, we re-prove and prove the following two results for ternary octics:

- for all nonnegative $f \in P_{\mathbb{P}^2,8}$, there exists a nonzero $g \in \Sigma_{\mathbb{P}^2,4}$ such that $fg \in \Sigma_{\mathbb{P}^2,12}$; and
- there exists nonnegative $f \in P_{\mathbb{P}^2,8}$ such that, for all nonzero $g \in \Sigma_{\mathbb{P}^2,2}$, we have $fg \notin \Sigma_{\mathbb{P}^2,10}$.

Together these give the first tight bounds on the degrees of sum-of-squares multipliers for homogeneous polynomials since Hilbert’s 1893 paper [Hil] in which he proves sharp bounds for ternary sextics. No other sharp bounds for homogeneous polynomials are known. For example, the recent theorem in [P] shows that, for quaternary quartics, one can multiply by sum-of-squares of degree 4 to obtain a sum of squares, but it is not known whether quadratic multipliers suffice.

By reinterpreting Theorem 1.1 or Theorem 1.2, we do obtain degree bounds for certificates of nonnegativity. A sum-of-squares multiplier g certifies that the element f is nonnegative at all points where g does not vanish. When the complement of this vanishing set is dense in the Euclidean topology, it follows that the element f is nonnegative. Changing perspectives, these theorems also generate a finite hierarchy of approximations to the cone $P_{X,2j}$, namely the sets $\{f \in R_{2j} : \text{there exists } g \in \Sigma_{X,2k} \text{ such that } fg \in \Sigma_{X,2j+2k}\}$; compare with Subsection 3.6.1 in [BPT]. It follows that deciding if an element f belongs to the cone $P_{X,2j}$ is determined by a semidefinite program of known size.

Relationship with prior results. Our degree bounds, with their uniformity and sharpness, cannot be directly compared to any established bound on multipliers, except for those on zero-dimensional schemes in [BGP]. Most earlier work focuses on general semi-algebraic sets, where no sharpness results are known, or on affine curves, where no uniform bounds are possible for singular curves.

The best bound on the degree of a sum-of-squares multiplier on an arbitrary semi-algebraic set involves a tower of five exponentials; see Theorem 1.5.7 in [LPR]. However, Corollary 4.9 shows that, for a nondegenerate totally-real projective curve $X \subset \mathbb{P}^n$ of degree d , every nonnegative form

admits a nonzero sum-of-squares multiplier of degree $2k$ for all $k \geq d - n + 1$. Absent sharp bounds in some larger context, it is impossible to ascertain if this difference in the complexity of the bounds is just a feature of low-dimensional varieties or part of some more general phenomenon.

Restricting to curves likewise fails to produce meaningful comparisons. Corollary 4.15 in [Sc1] illustrates that one can often certify nonnegativity without using a multiplier on an affine curve. Concentrating on a special type of multiplier, Theorem 4.11 in [Sc2] demonstrates that, on a nonsingular projective curve, any sufficiently large power of a positive element gives a multiplier; also see [R]. For nonsingular affine curves, Corollary 4.4 in [Sc3] establishes that there exist universal degree bounds, even though the techniques do not yield explicit bounds. In contrast with Theorem 1.1, the bounds in these situations depend on the nonnegative element f and tend towards positive infinity as the underlying curve acquires certain singularities.

To identify a close analogue of our work, we must lower the dimension: Theorems 1.1–1.2 in [BGP] provide uniform degree bounds over a finite set of points that are tight for quadratic functions on the hypercube. The lone additional sharp degree bound on multipliers is, to the best of our knowledge, Hilbert’s original work [Hil] on ternary sextics.

Main ideas. The results in this paper arose while exploring the relationship between convex geometry and algebraic geometry for sum of squares on real varieties. The two parts of our main theorems are proven independently. The upper bound on the minimum degree of a sum-of-squares multiplier is derived from a new Bertini Theorem in convex algebraic geometry and the lower bound is obtained by deforming rational Harnack curves on toric surfaces.

To prove the first part, we reinterpret the non-existence of a sum-of-square multiplier $g \in \Sigma_{X,2k}$ as asserting that the convex cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ intersect only at zero. If a real subscheme $X \subseteq \mathbb{P}^n$ possesses a linear functional separating these cones, then Theorem 3.1 demonstrates that a sufficiently general hypersurface section of X also does. In this setting, a phrase ‘sufficiently general’ means belonging to a nonempty open subset in the Euclidean topology of the relevant parameter space. Unexpectedly, this convex version of Bertini’s Theorem relies on our characterization of spectrahedral cones that have many facets in a neighbourhood of every point; see Proposition 2.5. Recognizing this dependency is the crucial insight. By repeated applications of our Bertini Theorem, we reduce to the case of points. Theorem 4.3 establishes the degree bound for the existence of a sum-of-squares multiplier on curves and Theorem 4.13 gives an higher-dimensional variant on arithmetically Cohen–Macaulay varieties.

To prove the second part, we show that having a nonnegative element vanish at a relatively large number of isolated real singularities precludes it from having a low-degree sum-of-squares multiplier. As Proposition 5.5 indicates, the hypotheses needed to actually realize this basic premise are formidable. Nonetheless, this transforms the problem into finding enough curves that satisfy the conditions and maximize the number of isolated real singularities. Proposition 5.7 confirms that rational singular Harnack curves on toric surfaces fulfill these requirements. By perturbing both the curve X and the nonnegative element $f \in R_{2j}$, Theorem 5.8 exhibits smooth curves and nonnegative elements without low-degree sum-of-squares multipliers. Proposition 5.15 then lifts these degree

bounds from curves to some surfaces. Miraculously, for totally-real projective curves, the degree bounds in Theorem 4.3 and Theorem 5.8 coincide.

Explicit Examples. Beyond the uniformity, our results also specialize to simple degree bounds in many interesting situations. As one might expect, the degree bounds are straightforward for complete intersections and planar curves; see Example 4.6 and Example 4.7. However, by demonstrating that our degree bound is sharp for some, but not all, planar curves, Example 4.8 and Example 5.3 are much more innovative. For certain non-planar curves lying on embedded toric surfaces, Examples 5.11–5.14 present sharp degree bounds. These examples also serve as our best justification for the second part of Theorem 1.1. It remains an open problem to classify all of the curves for which the bounds in Theorem 1.1 are sharp. Switching to higher-dimensional arithmetically Cohen–Macaulay varieties, Example 4.16 re-establishes that a nonnegative quadratic form on a totally-real variety of minimal degree is a sum of squares. Examples 4.17–4.19, which bound multipliers on surfaces of minimal degree, the projective plane, and surfaces of almost minimal degree respectively, appear to exhaust all of the consequential applications to surfaces. Together Example 4.17 and Example 5.16 establish Theorem 1.2. Highlighting the peculiarity of curves, this pair of examples also illustrate the gap between our upper and lower bounds on the minimal degree of a multiplier in this case. Nevertheless, Example 5.17 does give our new sharp bound for ternary octics. Despite being labelled examples, these are essential aspects of the paper.

2. MANY-FACETED SPECTRAHEDRAL CONES

This section focuses on convex geometry and properties of spectrahedral cones. We distinguish certain spectrahedral cones that have an abundance of facets in the vicinity of every point. To demonstrate the ubiquity of these cones in convex algebraic geometry, we show that if a sums-of-squares cone is closed and contains no lines, then its dual has this structure.

Let V be a finite-dimensional real vector space, let $S_2 := \text{Sym}^2(V^*)$ be the vector space of quadratic forms on V , and let $S_2^+ \subseteq S_2$ be the cone of positive-semidefinite quadratic forms. The corank of a quadratic form $f \in S_2$ is the dimension of $\text{Ker}(f)$, the kernel of the associated symmetric matrix. We endow $\text{Sym}^2(V^*)$ with the metric topology arising from the spectral norm. Since all norms on a finite-dimensional vector space induce the same topology, we refer to this metric topology as the Euclidean topology. For a quadratic form $g \in S_2$ and a positive real number ε , we write $B_\varepsilon(g) \subset S_2$ for the open ball of radius ε centered at g . The boundary ∂W of a subset $W \subseteq S_2$ equals the closure of W in the Euclidean topology without the interior of W . We also equip each subset $W \subseteq S_2$ with the induced Euclidean topology.

A linear subspace $L \subseteq S_2$ determines a spectrahedral cone $C := L \cap S_2^+$. The faces of the convex set C have a useful algebraic description. Specifically, Theorem 1 in [RG] establishes that the minimal face of C containing a given quadratic form $g \in S_2$ equals the intersection of C with the linear subspace consisting of $f \in S_2$ such that $\text{Ker}(g) \subseteq \text{Ker}(f)$. Hence, if the linear subspace L intersects the interior of the cone S_2^+ , then a quadratic form having corank 1 determines a facet, that is an inclusion-maximal proper face.

Our first lemma identifies a special type of spectrahedral cone. Given a nonzero $v \in V$, let $T_v \subset S_2$ denote the linear subspace consisting of the quadratic forms $f \in S_2$ such that $v \in \text{Ker}(f)$.

Lemma 2.1. *If the quadratic form $g \in \partial C$ has corank 1 and $\varepsilon > 0$ is sufficiently small, then the map $\varphi: B_\varepsilon(g) \cap \partial C \rightarrow \mathbb{P}(V)$, sending a quadratic form f to the linear subspace $\text{Ker}(f)$, is well-defined. Moreover, if the defining linear subspace L intersects the linear subspace $T_{\varphi(g)}$ transversely, then the image of φ contains a neighbourhood of $\varphi(g)$.*

Proof. The existence of $g \in \partial C$ having corank 1 implies that the defining linear subspace L meets the interior of S_2^+ . If not, then $C = L \cap S_2^+$ would be entirely contained in a face of S_2^+ , and all of its boundary points would have corank at least 2.

We claim that $\partial C = L \cap \partial S_2^+$. Since every neighbourhood of a point in ∂C contains at least one point in S_2^+ and at least one point not in S_2^+ , we have $\partial C \subseteq L \cap \partial S_2^+$. On the other hand, suppose that $f \in L \cap \partial S_2^+$ belongs to the relative interior of C . Since $f \in \partial S_2^+$, there exists a nonzero linear functional $\ell \in S_2^*$ that is nonnegative on S_2^+ and satisfies $\ell(f) = 0$. As f lies in the relative interior of C , it follows that ℓ vanishes identically on C and C is contained in $\ell^{-1}(0) \cap S_2^+ \subseteq \partial S_2^+$. However, this is absurd because L intersects the interior of S_2^+ , so we obtain $\partial C = L \cap \partial S_2^+$.

Since the eigenvalues of a matrix are continuous functions of the entries of the matrix, we see that, for a sufficiently small $\varepsilon > 0$, each point in $B_\varepsilon(g) \cap \partial S_2^+$ has corank 1. Hence, for each quadratic form $f \in B_\varepsilon(g) \cap \partial C$, the linear subspace $\text{Ker}(f)$ has dimension one. Therefore, the map $\varphi: B_\varepsilon(g) \cap \partial C \rightarrow \mathbb{P}(V)$ is well-defined.

To prove the second part of the lemma, let $m := \dim(V)$. By definition, we have $T_{\lambda v} = T_v$ for any nonzero $\lambda \in \mathbb{R}$. Moreover, requiring that a nonzero vector $v \in V$ belongs to the kernel of a symmetric $(m \times m)$ -matrix imposes m independent linear conditions on the entries of the matrix, so we have $\text{codim } T_v = m$. The hypothesis that the linear subspaces L and $T_{\varphi(g)}$ meet transversely means that $S_2 = L + T_{\varphi(g)}$, which implies that $\dim S_2 = \dim(L) + \dim(T_{\varphi(g)}) - \dim(L \cap T_{\varphi(g)})$ and $\dim(L \cap T_{\varphi(g)}) = \dim(L) - m$. It follows that, for all v in an open neighbourhood $W \subseteq \mathbb{P}(V)$ of $\varphi(g)$ in the Euclidean topology, we have $\dim(L \cap T_v) = \dim(L) - m$.

Consider the set $U_\varepsilon := \{[v] \in W : T_v \cap L \cap B_\varepsilon(g) \neq \emptyset\}$. Let $G := \text{Gr}(\dim(L) - m, L)$ be the Grassmannian of linear subspaces in L with codimension m considered as a real manifold, and let π_1, π_2 be the canonical projection maps from the universal family in $L \times G$ onto the factors. The previous paragraph shows that the map $\psi: W \rightarrow G$ sending $[v]$ to the linear subspace $L \cap T_v$ is well-defined and continuous. Since π_1 is a continuous map and π_2 is an open map, we see that $U_\varepsilon = \psi^{-1}(\pi_2(\pi_1^{-1}(L \cap B_\varepsilon(g))))$ is an open subset in W considered as a real manifold.

Finally, if the image of φ does not contain U_ε for all sufficiently small $\varepsilon > 0$, then there is a sequence of increasing positive integers r_i , a sequence of nonzero vectors $v_i \in V$, and a sequence of quadratic forms $f_i \in S_2 \setminus S_2^+$ such that the $[v_i] \in \mathbb{P}(V)$ converge to $\varphi(g)$ as $i \rightarrow \infty$ and $f_i \in L \cap T_{v_i} \cap B_{1/r_i}(g)$ for each $i \in \mathbb{N}$. Thus, the quadratic forms f_i converge to g as $i \rightarrow \infty$. However, for sufficiently large $i \in \mathbb{N}$, the symmetric matrix corresponding to f_i has a negative eigenvalue and corank 1, because $f_i \in S_2 \setminus S_2^+$ and $f_i \in B_{1/r_i}(g)$. Hence, the limit of the f_i cannot be both positive-semidefinite and have corank 1. We conclude that, for sufficiently small ε , the elements in

$L \cap T_V \cap B_\varepsilon(g)$ for all $[v] \in U_\varepsilon$ are positive-semidefinite. Therefore, the image of φ contains U_ε for sufficiently small ε . \square

Building on Lemma 2.1, we introduce the following class of spectrahedral cones. This definition guarantees that, both globally and locally, the spectrahedral cone has numerous facets.

Definition 2.2. A spectrahedral cone $C = L \cap S_2^+$ is *many-faceted* if the points with corank 1 form a dense subset of ∂C and, for all $g \in \partial C$ with corank 1 and all sufficiently small $\varepsilon > 0$, the image of $\varphi: B_\varepsilon(g) \cap \partial C \rightarrow \mathbb{P}(V)$ contains a neighbourhood of $\varphi(g)$.

Being many-faceted is an extrinsic property; it depends on the presentation of the spectrahedral cone. For instance, if $C = L \cap S_2^+$ is many-faceted, then we must have $\dim(L) \geq \dim(V) + 1$.

Two modest examples help illuminate this definition.

Example 2.3 (A spectrahedral cone that is not many-faceted). Let $V = \mathbb{R}^3$ and $S_2 = \mathbb{R}[x_0, x_1, x_2]_2$. For the spectrahedral cone $C := L \cap S_2^+$ given by the linear subspace $L := \text{Span}(x_0^2, x_0x_2, x_1^2, x_2^2) \subset S_2$, we have $C = \{\alpha x_0^2 + 2\beta x_0x_2 + \gamma x_1^2 + \delta x_2^2 : \alpha \geq 0, \gamma \geq 0, \text{ and } \alpha\delta - \beta^2 \geq 0\}$ and the associated symmetric matrices have the form $\begin{bmatrix} \alpha & 0 & \beta \\ 0 & \gamma & 0 \\ \beta & 0 & \delta \end{bmatrix}$. The relative interior of the face given by $\gamma = 0$ is open in ∂C and consists of points with corank 1, because the kernel of each quadratic form in the relative interior of this face is equal to $\text{Span}([0 \ 1 \ 0]^T)$. However, the image of the map φ is a single point in $\mathbb{P}(V)$, so it does not contain an open subset. Thus, this spectrahedral cone is not many-faceted. \diamond

Example 2.4 (A spectrahedral cone that is many-faceted). As in Example 2.3, let $V = \mathbb{R}^3$ and $S_2 = \mathbb{R}[x_0, x_1, x_2]_2$. For $C := L \cap S_2^+$ defined by $L := \text{Span}(x_0^2 + x_1^2, x_0^2 + x_2^2, x_0x_1, x_0x_2, x_1x_2) \subset S_2$, it follows that

$$C = \left\{ \alpha x_0^2 + 2\beta x_0x_1 + 2\gamma x_0x_2 + \alpha x_1^2 + 2\delta x_1x_2 + \alpha x_2^2 : \begin{array}{l} \alpha \geq 0, \alpha^2 - \beta^2 \geq 0, \\ \alpha^2 - \delta^2 \geq 0, \alpha^2 - \gamma^2 \geq 0, \text{ and} \\ \alpha^3 - \alpha\beta^2 - \alpha\gamma^2 - \alpha\delta^2 + 2\beta\gamma\delta \geq 0 \end{array} \right\}$$

and the associated symmetric matrices have the form $\begin{bmatrix} \alpha & \beta & \gamma \\ \beta & \alpha & \delta \\ \gamma & \delta & \alpha \end{bmatrix}$. The algebraic boundary of the section of this cone determined by setting $\alpha = 1$ equals the Cayley cubic surface defined by affine equation $1 - \beta^2 - \gamma^2 - \delta^2 + 2\beta\gamma\delta$; see Subsection 5.2.2 in [BPT]. From the well-known image of the boundary surface (see Figure 2.4.1), which is affectionately referred to as ‘The Samosa’, we see that the cone is many-faceted. For the quadratic form $g := x_0^2 + (x_1 - x_2)^2 \in \partial C$, we have $\varphi(g) = [0 : 1 : 1] \in \mathbb{P}(V)$ and

$$\dim(L \cap T_{\varphi(g)}) = \dim \text{Span}(x_1^2 - x_2^2, x_0x_1, x_0x_2) = 3 > 2 = \dim(L) - \dim(V).$$

This shows that there exist many-faceted spectrahedral cones which do not arise via Lemma 2.1. \diamond

To realize such many-faceted cones within convex algebraic geometry, consider a real projective subscheme $X \subseteq \mathbb{P}^n = \text{Proj}(S)$ where $S := \mathbb{R}[x_0, x_1, \dots, x_n]$. If I_X is the saturated homogeneous



FIGURE 2.4.1. Two perspectives of ‘The Samosa’

ideal defining X , then the \mathbb{Z} -graded coordinate ring of X is $R := S/I_X$. For each $j \in \mathbb{Z}$, the graded component R_j of degree j is a finite-dimensional real vector space, and we set

$$\Sigma_{X,2j} := \{f \in R_{2j} : \text{there exists } g_0, g_1, \dots, g_s \in R_j \text{ such that } f = g_0^2 + g_1^2 + \dots + g_s^2\}.$$

Since a nonnegative real number has a square root in \mathbb{R} , we see that $\Sigma_{X,2j}$ is a convex cone in R_{2j} . The map $\sigma_j: \text{Sym}^2(R_j) \rightarrow R_{2j}$, induced by multiplication, is surjective. It follows that the cone $\Sigma_{X,2j}$ is also full-dimensional, because the second Veronese embedding of \mathbb{P}^n is nondegenerate. Moreover, the dual map $\sigma_j^*: R_{2j}^* \rightarrow \text{Sym}^2(R_j^*)$ is injective and, for $\ell \in R_{2j}^*$, the symmetric form $\sigma_j^*(\ell): R_j \otimes_{\mathbb{R}} R_j \rightarrow \mathbb{R}$ is given explicitly by $g_1 \otimes g_2 \mapsto \ell(g_1 g_2)$.

The subsequent proposition consolidates a few fundamental properties of this cone and proves that many-faceted spectrahedral cones are common in convex algebraic geometry. A cone in a real vector space is **pointed** if it is both closed in the Euclidean topology and contains no lines.

Proposition 2.5. *Fix $j \in \mathbb{N}$. If $X \subseteq \mathbb{P}^n$ is a real projective subscheme with \mathbb{Z} -graded coordinate ring R such that the map $\eta_g: R_j \rightarrow R_{2j}$ defined by $\eta_g(f) = fg$ is injective for all nonzero $g \in R_j$, then the following are equivalent.*

- (a) *The cone $\Sigma_{X,2j}$ is pointed.*
- (b) *No nontrivial sum of squares of forms of degree j equals zero.*
- (c) *The points of corank 1 form a dense subset of $\partial \Sigma_{X,2j}^*$ in the Euclidean topology.*
- (d) *The dual $\Sigma_{X,2j}^*$ is a many-faceted spectrahedral cone.*

Proof. (a) \Rightarrow (b): If some nontrivial sum of squares equals zero, then there exist $g_0, g_1, \dots, g_s \in R_j$ satisfying $g_0^2 + g_1^2 + \dots + g_s^2 = 0$. We have $s > 0$, because the map η_g is injective for all nonzero $g \in R_j$. Since $g_0^2 = -(g_1^2 + g_2^2 + \dots + g_s^2)$, it follows that $\lambda g_0^2 \in \Sigma_{X,2j}$ for all $\lambda \in \mathbb{R}$ which contradicts the assumption that $\Sigma_{X,2j}$ contains no lines.

(b) \Rightarrow (a): Fix an inner product on the real vector space R_j and let $g \mapsto \|g\|$ denote the associated norm. The spherical section $K := \{g^2 \in R_{2j} : g \in R_j \text{ satisfies } \|g\| = 1\}$ is compact, because it is the continuous image of a compact set. Moreover, the convex hull of K does not contain 0, because no nontrivial sum of squares equals zero. Since $\Sigma_{X,2j}$ is the conical hull of K , the cone $\Sigma_{X,2j}$ is closed. If $\Sigma_{X,2j}$ contains a line, then there exists a nonzero $f \in R_{2j}$ such that both f and $-f$ lie in $\Sigma_{X,2j}$. However, it follows that the nontrivial sum $f + (-f)$ equals zero, which contradicts (b).

(a) \Rightarrow (c): Since $\Sigma_{X,2j}$ is a pointed full-dimensional cone, its dual $\Sigma_{X,2j}^*$ is also a pointed full-dimensional cone. As a consequence, Theorem 2.2.4 in [ScR] implies that the linear functionals

$\ell \in \Sigma_{X,2j}^*$ whose normal cone is a single ray form a dense subset of $\partial \Sigma_{X,2j}^*$. We claim that every such linear functional ℓ has corank one. If $f, g \in R_j$ are two nonzero elements lying in the kernel of $\sigma_j^*(\ell)$, then f^2 and g^2 are nonzero elements of the normal cone of $\Sigma_{X,2j}^*$ at ℓ . Because this normal cone is a ray, there exists a positive $\lambda \in \mathbb{R}$ such that $f^2 = \lambda g^2$. Hence, we have $(f + \sqrt{\lambda}g)(f - \sqrt{\lambda}g) = 0$ in R_{2j} . By injectivity of multiplication maps, we conclude that f and g are linearly dependent, so ℓ has corank 1 and (c) holds.

(c) \Rightarrow (a): If $\Sigma_{X,2j}$ is not closed, then the ‘(b) \Rightarrow (a)’ step shows that there is a nontrivial sum of squares from R_j equal to zero in R_{2j} . In this case, the ‘(a) \Rightarrow (b)’ step demonstrates that $\Sigma_{X,2j}$ contains a line. Now, if $\Sigma_{X,2j}$ contains a line, then its dual $\Sigma_{X,2j}^*$ is not full-dimensional. Since the dual map $\sigma_j^*: R_{2j}^+ \rightarrow \text{Sym}^2(R_j^*)$ is injective, the linear subspace $\sigma_j^*(R_{2j}^+)$ does not intersect the interior of the cone S_2^+ consisting of positive-semidefinite forms in $\text{Sym}^2(R_j^*)$. Hence, the image $\sigma_j^*(\Sigma_{X,2j}^*)$ consists of symmetric forms of corank at least 1. It follows that the boundary consists of symmetric forms of corank at least 2, which contradicts (c).

(a) \Leftrightarrow (d): Let $V := R_j$ and let $S_2 := \text{Sym}^2(R_j^*)$. For a linear functional $\ell \in \Sigma_{X,2j}^*$, we have $\ell(g^2) \geq 0$ for all $g \in R_j$, so the symmetric form $\sigma_j^*(\ell)$ is positive-semidefinite. Conversely, if $\sigma_j^*(\ell)$ is positive-semidefinite symmetric form, then we have $\ell(g^2) \geq 0$ for all $g \in R_j$. It follows that $\ell(g_0^2 + g_1^2 + \dots + g_s^2) = \ell(g_0^2) + \ell(g_1^2) + \dots + \ell(g_s^2) \geq 0$ for $g_0, g_1, \dots, g_s \in R_j$ and $\ell \in \Sigma_{X,2j}^*$. Hence, the map σ_j^* identifies the dual $\Sigma_{X,2j}^*$ with the spectrahedral cone determined by the linear subspace $L := \sigma_j^*(R_{2j}^+)$ in $S_2 = \text{Sym}^2(R_j^*)$; compare with Lemma 2.1 in [BSV]. Given a nonzero $f \in V$, let $T_f \subset S_2$ be the linear subspace consisting of the symmetric forms $h \in S_2$ such that $f \in \text{Ker}(h)$. As in the proof of Lemma 2.1, we have $\text{codim } T_f = \dim V$. The map σ_j^* identifies the linear subspace $L \cap T_f$ with the set of linear functionals $\ell \in R_{2j}^*$ such that $\ell(fg) = 0$ for all $g \in V$. If $\langle f \rangle$ denotes the ideal in R generated by f , then it follows that the codimension of $L \cap T_f$ in L equals the dimension of $\langle f \rangle_{2j}$. By hypothesis, the map $\eta_f: R_j \rightarrow R_{2j}$ is injective, so we obtain $\dim \langle f \rangle_{2j} = \dim R_j = \dim V$. Hence, we have $\dim L + \dim T_f - \dim L \cap T_f = \dim S_2$ and the linear subspaces L and T_f meet transversely for all nonzero $f \in V$. If $g \in \partial \Sigma_{X,2j}^*$ has corank 1 and $\varepsilon > 0$ is sufficiently small, then Lemma 2.1 establishes that the image of $\varphi: B_\varepsilon(g) \cap \partial \Sigma_{X,2j}^* \rightarrow \mathbb{P}(V)$ contains a neighbourhood of $\varphi(g)$. Since ‘(a) \Leftrightarrow (c)’ establishes that $\Sigma_{X,2j}$ is pointed if and only if the points of corank 1 form a dense subset of $\partial \Sigma_{X,2j}^*$ in the Euclidean topology, we conclude that $\Sigma_{X,2j}$ is pointed if and only if its dual $\Sigma_{X,2j}^*$ is a many-faceted spectrahedral cone. \square

Remark 2.6. The first condition in Proposition 2.5 may be rephrased using different terminology. A cone C is salient if it does not contain an opposite pair of nonzero vectors, that is $(-C) \cap C \subseteq \{0\}$. In other words, a cone is salient if and only if it contains no lines. Hence, a cone is pointed if it is both closed and salient.

Remark 2.7. If $\Sigma_{X,2j}$ is not closed, then the ‘(c) \Rightarrow (a)’ proves that $\Sigma_{X,2j}$ contains a line.

We end this section with special cases of Proposition 2.5. A subscheme $X \subseteq \mathbb{P}^n$ is a real projective variety if it is a geometrically integral projective scheme over \mathbb{R} . Moreover, a real variety X is totally

real if the set $X(\mathbb{R})$ of real points is Zariski dense. The most important application of Proposition 2.5 is the following corollary.

Corollary 2.8. *If $X \subseteq \mathbb{P}^n$ is a real projective variety, then the cone $\Sigma_{X,2j}$ is pointed if and only if its dual $\Sigma_{X,2j}^*$ is a many-faceted spectrahedral cone. Furthermore, the cones $\Sigma_{X,2j}^*$ are many-faceted for all $j \in \mathbb{N}$ if and only if X is totally real.*

Proof. Because X is geometrically integral, its coordinate ring R is a domain. Hence, each nonzero element in R_j is a nonzerodivisor and the map $\eta_g : R_j \rightarrow R_{2j}$ is injective for all nonzero $g \in R_j$. By combining this observation with Proposition 2.5, we first conclude that $\Sigma_{X,2j}$ is pointed if and only if its dual $\Sigma_{X,2j}^*$ is a many-faceted spectrahedral cone. Secondly, X is totally-real if and only if, for all $j \in \mathbb{N}$, no nontrivial sum of squares from R_j equals zero in R_{2j} . Therefore, the first part together with Proposition 2.5 establishes that the cones $\Sigma_{X,2j}^*$ are many-faceted for every $j \in \mathbb{N}$ if and only if X is totally real. \square

3. A BERTINI THEOREM FOR SEPARATORS

In this section, we explore the properties of separating hyperplanes within convex algebraic geometry. Two cones C_1 and C_2 in a real vector space are **well-separated** if there exists a linear functional ℓ such that $\ell(v) > 0$ for all nonzero $v \in C_1$ and $\ell(v) < 0$ for all nonzero $v \in C_2$. A linear functional ℓ with these properties is called a **strict separator**. If C_1 and C_2 are pointed, then being well-separated is equivalent to $C_1 \cap C_2 = \{0\}$.

The main result in this section is an analogue of Bertini's Theorem to convex algebraic geometry. As in Section 2, $X \subseteq \mathbb{P}^n$ is a real projective subscheme with \mathbb{Z} -graded coordinate ring $R = S/I_X$ and $S := \mathbb{R}[x_0, x_1, \dots, x_n]$. Given an element $f \in R_{2j}$, we set $f \cdot \Sigma_{X,2k} := \{fg \in R_{2j+2k} : g \in \Sigma_{X,2k}\}$. For a nonzero homogeneous polynomial $h \in S$, the associated hypersurface section of X is the subscheme $X' := X \cap V(h) \subset \mathbb{P}^n$. The \mathbb{Z} -graded coordinate ring of X' is the quotient $R' := S/I_{X'}$ where $I_{X'}$ is the saturated homogeneous ideal $(I_X + \langle h \rangle : \langle x_0, x_1, \dots, x_n \rangle^\infty)$. We write $f' \in R'_{2j}$ for the canonical image of $f \in R_{2j}$.

Theorem 3.1. *Fix positive integers j and k . Let $X \subseteq \mathbb{P}^n$ be a real projective subscheme with coordinate ring R such that the map $\eta_g : R_{j+k} \rightarrow R_{2j+2k}$ is injective for all nonzero $g \in R_{j+k}$, and consider a nonzerodivisor $f \in R_{2j}$. If the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated, then the set of hypersurface sections X' of X , such that $\Sigma_{X',2j+2k}$ and $f' \cdot \Sigma_{X',2k}$ are well-separated, contains a nonempty open subset of $\mathbb{P}(R_{j+k})$ in the Euclidean topology.*

Proof. To begin, we prove that the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are pointed. By hypothesis, $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated, so neither cone contains a line. Hence, Remark 2.7 shows that $\Sigma_{X,2j+2k}$ is also closed. As f is a nonzerodivisor, the map $\eta_f : R_{2k} \rightarrow R_{2j+2k}$ is injective, so the cone $\Sigma_{X,2k}$ is isomorphic to the cone $f \cdot \Sigma_{X,2k}$. Hence, a second application of Remark 2.7 establishes that both $\Sigma_{X,2k}$ and $f \cdot \Sigma_{X,2k}$ are closed.

Now, let $C := \Sigma_{X,2j+2k}^* \cap (-f \cdot \Sigma_{X,2k})^*$ be the cone of separators. The cone C is closed and full-dimensional, because $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. In particular, the boundary of

C is not contained in the boundary of $\Sigma_{X,2j+2k}^*$ or the boundary of $(-f \cdot \Sigma_{X,2k})^*$. Since the cone $(-f \cdot \Sigma_{X,2k})^*$ is full-dimensional and

$$\begin{aligned} D &:= \partial C \setminus \partial(-f \cdot \Sigma_{X,2k})^* \\ &= \left((\partial(-f \cdot \Sigma_{X,2k})^* \cap \Sigma_{X,2j+2k}^*) \cup ((-f \cdot \Sigma_{X,2k})^* \cap \partial \Sigma_{X,2j+2k}^*) \right) \setminus \partial(-f \cdot \Sigma_{X,2k})^* \\ &= ((-f \cdot \Sigma_{X,2k})^* \setminus \partial(-f \cdot \Sigma_{X,2k})^*) \cap \partial \Sigma_{X,2j+2k}^*, \end{aligned}$$

it follows that D is a nonempty open subset of $\partial \Sigma_{X,2j+2k}^*$ in the Euclidean topology. Since $\Sigma_{X,2j+2k}$ is pointed, Proposition 2.5 implies that $\Sigma_{X,2j+2k}^*$ is a many-faceted spectrahedral cone. Hence, the points with corank 1 form a dense subset of $\partial \Sigma_{X,2j+2k}^*$, so we may choose $g \in D$ with corank 1. Moreover, for a sufficiently small $\varepsilon > 0$, the image of the map $\varphi: B_\varepsilon(g) \cap D \rightarrow \mathbb{P}(R_{j+k})$ contains a neighbourhood U of $\varphi(g)$. Hence, if $\ell \in R_{2j+2k}^*$ satisfies $[\varphi(\ell)] \in U$, then there exists $h \in R_{j+k}$ such that $\text{Ker } \sigma_{j+k}^*(\ell) = \text{Span}(h)$. Let $X' := X \cap V(h)$ denote the corresponding hypersurface section with coordinate ring R' . Since ℓ' has corank 1, the linear functional $\ell \in R_{2j+2k}^*$ induces a strict separator $\ell' \in (R')_{2j+2k}^*$ on the cones $\Sigma_{X',2j+2k}$ and $f' \cdot \Sigma_{X',2k}$. Therefore, the set of X' , such that $\Sigma_{X',2j+2k}$ and $f' \cdot \Sigma_{X',2k}$ are well-separated, contains the nonempty open subset U of $\mathbb{P}(R_{j+k})$. \square

To exploit Theorem 3.1, we also need to understand the properties of strict separators on zero-dimensional schemes. As we will see, the existence of strict separators imposes nontrivial constraints on a set of points. For a real projective scheme $X \subseteq \mathbb{P}^n$ with homogeneous coordinate ring R , the Hilbert function $h_X: \mathbb{Z} \rightarrow \mathbb{Z}$ is defined by $h_X(j) := \dim_{\mathbb{R}} R_j$. Following Section 3.1 in [Ha3] or Section 2 in [Ha1], a set of points $X \subseteq \mathbb{P}^n$, that is a zero-dimensional reduced subscheme, has the **uniform position property** if the Hilbert function of a subset of X depends only on the cardinality of the subset.

The concluding proposition of this section shows that the existence of certain positive linear functionals on a set of points imposes constraints on its Hilbert function.

Proposition 3.2. *Fix positive integers j and k , and let $X \subseteq \mathbb{P}^n$ be a set of at least two points with the uniform position property.*

- (i) *Suppose that X has no real points. If $\ell \in R_{2k}^*$ is positive on the nonzero elements in $\Sigma_{X,2k}$, then we have $h_X(k) \leq \lceil \frac{1}{2} h_X(2k) \rceil$.*
- (ii) *Suppose that $f \in R_{2j}$ is positive on $X(\mathbb{R})$ and doesn't vanish at any point in $X(\mathbb{C})$. If $\ell \in R_{2j+2k}^*$ is a strict separator for $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$, then we have $h_X(k) + h_X(j+k) \leq h_X(2j+2k)$.*

Proof. We begin with an analysis of the symmetric forms arising from point evaluations. Let $Z \subseteq X$ be a subset consisting of e distinct real points and m complex conjugate pairs. Choose affine representatives $\tilde{p}_1, \tilde{p}_2, \dots, \tilde{p}_e \in \mathbb{A}^{n+1}(\mathbb{R})$ for points in $Z(\mathbb{R})$, and choose affine representatives $\tilde{a}_1 \pm \tilde{b}_1 \sqrt{-1}, \tilde{a}_2 \pm \tilde{b}_2 \sqrt{-1}, \dots, \tilde{a}_m \pm \tilde{b}_m \sqrt{-1} \in \mathbb{A}^{n+1}(\mathbb{R})$ where $\tilde{a}_j, \tilde{b}_j \in \mathbb{A}^{n+1}(\mathbb{R})$ for the complex conjugate pairs in $Z(\mathbb{C})$. For $p \in X(\mathbb{R})$ and $k \in \mathbb{N}$, evaluation at an affine representative $\tilde{p} \in \mathbb{A}^{n+1}(\mathbb{R})$ determines the linear functional $\tilde{p}^* \in R_{2k}^*$. Any linear functional $\ell \in R_{2k}^*$ lying in the span of these

point evaluations can be written as

$$\ell = \sum_{i=1}^e \kappa_i \tilde{p}_i^* + \sum_{j=1}^m ((\lambda_j + \mu_j \sqrt{-1})(\tilde{a}_j + \tilde{b}_j \sqrt{-1})^* + (\lambda_j - \mu_j \sqrt{-1})(\tilde{a}_j - \tilde{b}_j \sqrt{-1})^*)$$

where $\kappa_i, \lambda_j, \mu_j \in \mathbb{R}$ for $1 \leq i \leq e$ and $1 \leq j \leq m$. It follows that

$$\sigma_{2k}^*(\ell) = \sum_{i=1}^e \kappa_i (\tilde{p}_i^*)^2 + \sum_{j=1}^m \lambda_j ((\tilde{a}_j^*)^2 - (\tilde{b}_j^*)^2) - 2\mu_j (\tilde{a}_j^*)(\tilde{b}_j^*) \in \text{Sym}^2(R_k^*).$$

The eigenvalues for the symmetric matrix $\begin{bmatrix} \lambda_j & -\mu_j \\ -\mu_j & -\lambda_j \end{bmatrix}$ are $\pm \sqrt{\lambda_j^2 + \mu_j^2}$, so the number of positive eigenvalues for $\sigma_{2k}^*(\ell)$ equals the number e_+ of positive κ_i plus the number m' of nonzero $\lambda_j^2 + \mu_j^2$. Similarly, the number of negative eigenvalues for $\sigma_{2k}^*(\ell)$ equals the number e_- of negative κ_i plus the number m' of nonzero $\lambda_j^2 + \mu_j^2$. Hence, if $\sigma_{2k}^*(\ell)$ is positive-definite, then we have $\text{h}_X(k) = \dim R_k^* \leq e_+ + m'$.

Using this analysis, we prove (i). Assume that $\ell \in R_{2k}^*$ is positive on the nonzero elements in $\Sigma_{X,2k}$. A form in R_{2k}^* is zero if and only if it is annihilated by $\tilde{p}^* \in R_{2k}^*$ for all points $p \in X(\mathbb{C})$. Hence, every linear functional in R_{2k}^* can be written as a \mathbb{C} -linear combinations of such point evaluations. The evaluations at the points in any subset X , with cardinality at least $\text{h}_X(2k)$, span R_{2k}^* , because X has the uniform position property. As X is a set of points, the value of Hilbert function $\text{h}_X(k)$ is at most the number of points. Since $X(\mathbb{R}) = \emptyset$, we may choose m conjugate pairs of points in $X(\mathbb{C})$ with $m := \lceil \frac{1}{2} \text{h}_X(2k) \rceil$; in particular, we have $e = 0$. Since $\sigma_{2k}^*(\ell)$ is positive-definite, the first paragraph shows that $\text{h}_X(k) \leq m' \leq m = \lceil \frac{1}{2} \text{h}_X(2k) \rceil$ as required.

We next examine the symmetric forms induced by the element $f \in R_{2k}$. Given $\ell \in R_{2k+2k}^*$, the linear functional $\ell' \in R_{2k}^*$ is defined by $\ell'(g) := \ell(fg)$ for all $g \in R_{2k}$. When $\ell \in R_{2j+2k}^*$ lies in the span of the point evaluations for Z , the expression for ℓ' as a linear combination of the point evaluations has the same number of positive, negative, and nonzero coefficients as ℓ , because $f \in R_{2j}$ is positive on $X(\mathbb{R})$ and does not vanish at any points in $X(\mathbb{C})$. Hence, if $\sigma_{2k}^*(\ell')$ is negative-definite, then the first paragraph implies that $\text{h}_X(k) = \dim R_k^* \leq e_- + m'$.

Lastly, we establish (ii). Assume that $\ell \in R_{2j+2k}^*$ is a strict separator for $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$. As in the second paragraph, we may choose a subset of X such that the point evaluations span R_{2j+2k}^* . Suppose that there exists a conjugate-invariant basis of R_{2j+2k}^* consisting of point evaluations at e distinct real points and m complex conjugate pairs. Since $\sigma_{2k}^*(\ell')$ is negative-definite and $\sigma_{2j+2k}^*(\ell)$ is positive-definite, the first and third paragraphs combine to show that

$$\text{h}_X(k) + \text{h}_X(j+k) \leq (e_- + m') + (e_+ + m') \leq e + 2m = \text{h}_X(2j+2k).$$

On the other hand, if no subset of X yields a conjugate-invariant basis of R_{2j+2k}^* , then there are $m := \lceil \frac{1}{2} \text{h}_X(2j+2k) \rceil$ conjugate pairs of points in $X(\mathbb{C})$ that span R_{2j+2k}^* and we have $e = 0$. Hence, we obtain $\text{h}_X(j+k) \leq m$, $2\text{h}_X(j+k) \leq 2m = \text{h}_X(2j+2k) + 1$, and $2\text{h}_X(j+k) - 1 \leq \text{h}_X(2j+2k)$. With the goal of finding a contradiction, assume that $\text{h}_X(k) + \text{h}_X(j+k) > \text{h}_X(2j+2k)$. It follows that $\text{h}_X(k) + 1 > \text{h}_X(j+k)$. The Hilbert function of a set of points is strictly increasing until it stabilizes

at the number points, so we deduce that $h_X(k) = h_X(j+k) = h_X(2j+2k)$. Hence, the inequality $2h_X(j+k) \leq h_X(2j+2k) + 1$ implies that $h_X(j+k) = 1$. However, this contradicts the hypothesis that X has at least two points. Therefore, we conclude that $h_X(k) + h_X(j+k) \leq h_X(2j+2k)$. \square

4. UPPER BOUNDS FOR SUM-OF-SQUARES MULTIPLIERS

This section establishes an upper bound on the minimal degree of a sum-of-squares multiplier. These geometric degree bounds for the existence of multipliers prove the first half of our main theorems. After a preparatory lemma, Theorem 4.3 describes the general result for curves and is followed by several corollaries and valuable examples. The same approach is then applied to higher dimensional varieties to obtain the general Theorem 4.13. The ensuing examples illustrate the applicability of this theorem.

Throughout this section, we work with a real projective subscheme $X \subseteq \mathbb{P}^n$ with \mathbb{Z} -graded coordinate ring $R = S/I_X$. The sign of an element $f \in R_{2j}$ at a real point $p \in X(\mathbb{R})$ is defined to be $\text{sgn}_p(f) := \text{sgn}(\tilde{f}(\tilde{p})) \in \{-1, 0, 1\}$, where the polynomial $\tilde{f} \in S_{2j}$ maps to f and the nonzero real point $\tilde{p} \in \mathbb{A}^{n+1}(\mathbb{R})$ maps to p under the canonical quotient maps. Since $p \in X(\mathbb{R})$, the real number $\tilde{f}(\tilde{p})$ is independent of the choice of \tilde{f} . Similarly, the choice of affine representative \tilde{p} is determined up to a nonzero real number and the degree of f is even, so the value of $\tilde{f}(\tilde{p})$ is determined up to the square of a nonzero real number. Hence, the sign of $f \in R_{2j}$ at $p \in X(\mathbb{R})$ is well-defined. We simply write $f(p) \geq 0$ for $\text{sgn}_p(f) \geq 0$. The subset $P_{X,2j} := \{f \in R_{2j} : f(p) \geq 0 \text{ for all } p \in X(\mathbb{R})\}$ forms a pointed full-dimensional convex cone in R_{2j} ; see Lemma 2.1 in [BSV].

As our initial focus, a **curve** $X \subseteq \mathbb{P}^n$ is a one-dimensional projective variety. Following Chapter 2 of [M], the **deficiency module** (also known as the Hartshorne-Rao module) of X is the \mathbb{Z} -graded S -module $M_X := \bigoplus_{i \in \mathbb{Z}} H^1(\mathbb{P}^n, \mathcal{I}_X(i))$. A homogeneous polynomial $h \in S = \mathbb{R}[x_0, x_1, \dots, x_n]$ determines the \mathbb{Z} -graded submodule $\text{Ann}_{M_X}(h) := (0 :_{M(X)} h) = \{f \in M_X : fh = 0\}$ of the deficiency module M_X . The next lemma (cf. Proposition 2.1.2 in [M]) shows that this submodule measures the failure of the ideal $I_X + \langle h \rangle$ to be saturated.

Lemma 4.1. *Fix a positive integer j and a nonnegative integer k . Let $X \subseteq \mathbb{P}^n$ be a curve. If $h \in S_k$ does not belong to the ideal I_X and $X' := X \cap V(h)$ is the associated hypersurface section of X , then we have $\text{Ann}_{M_X}(h)_{j-k} = 0$ if and only if $(I_{X'})_j = (I_X + \langle h \rangle)_j$.*

Proof. By definition, the submodule $\text{Ann}_{M_X}(h)$ fits into the exact sequence

$$0 \longrightarrow (\text{Ann}_{M_X}(h))(-k) \longrightarrow M_X(-k) \xrightarrow{\cdot h} M_X.$$

Sheafifying the canonical short exact sequence $0 \longrightarrow I_X \cap \langle h \rangle \longrightarrow I_X \oplus \langle h \rangle \longrightarrow I_X + \langle h \rangle \longrightarrow 0$ and taking cohomology of appropriate twists produces the long exact sequence

$$0 \longrightarrow I_X(-k) \xrightarrow{\begin{bmatrix} h \\ h \end{bmatrix}} I_X \oplus \langle h \rangle \longrightarrow I_{X'} \longrightarrow M_X(-k) \xrightarrow{\cdot h} M_X.$$

Breaking this long exact sequence into short exact sequences, we obtain

$$0 \longrightarrow I_X + \langle h \rangle \longrightarrow I_{X'} \longrightarrow (\text{Ann}_{M_X}(h))(-k) \longrightarrow 0.$$

Thus, we have $(\text{Ann}_{M_X}(h))(-k) \cong I_{X'}/(I_X + \langle h \rangle)$ and the required equivalence follows. \square

As a consequence of Lemma 4.1, we see that some natural geometric conditions imply that the ideal $I_X + \langle h \rangle$ is saturated.

Remark 4.2. A curve X is projectively normal if and only if $M_X = 0$. With this hypothesis, Lemma 4.1 implies that we have $(I_{X'})_j = (I_X + \langle h \rangle)_j$ for all $j \in \mathbb{Z}$. In particular, if X is arithmetically Cohen–Macaulay, then the ideal $I_X + \langle h \rangle$ is saturated.

The next result is the general form of our degree bound for the existence of sum-of-squares multipliers on curves.

Theorem 4.3. *Fix a positive integer j and a nonnegative integer k . Let $X \subseteq \mathbb{P}^n$ be a totally-real curve such that $H^1(\mathbb{P}^n, \mathcal{I}_X(j+k)) = 0$ and $h_X(2j+2k) < 2h_X(j+k) + h_X(k) - 1$. If $f \in P_{X,2j}$, then there exists a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$.*

Proof. We start by reinterpreting the non-existence of a suitable $g \in \Sigma_{X,2k}$ as the existence of a strict separator between appropriate cones. Corollary 2.8 implies that the cones $\Sigma_{X,2j+2k}$ and $\Sigma_{X,2k}$ are pointed. If $f = 0$, then the conclusion is trivial, so we may assume that f is nonzero. It follows that f is a nonzerodivisor, because X is integral. Since the map $\eta_f: R_{2k} \rightarrow R_{2j+2k}$ is injective, the pointed cone $\Sigma_{X,2k}$ is isomorphic to the cone $f \cdot \Sigma_{X,2k}$. Hence, the non-existence of a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$ is equivalent to saying that the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated.

To complete the proof, we reduce to the case of points by using new and old Bertini Theorems. Our convex variant, Theorem 3.1, implies that the set of homogeneous polynomials $h \in S_{j+k}$, such that $h \notin I_X$, $X' := X \cap V(h) \subset \mathbb{P}^n$, and the cones $\Sigma_{X',2j+2k}$ and $f' \cdot \Sigma_{X',2k}$ are well-separated, contains a nonempty Euclidean open subset $U_1 \subseteq \mathbb{P}(R_{j+k})$. Bertini's Theorem (see Théorème 6.3 in [J]) shows that there is a nonempty Zariski open subset $U_2 \subseteq \mathbb{P}(R_{j+k})$ such that, for all $[h] \in U_2$, the hypersurface section X' is a reduced set of points and f does not vanish at any point in X' . Moreover, our hypothesis that $(M_X)_{j+k} = 0$ combined with Lemma 4.1 establishes that there exists another nonempty Zariski open subset $U_3 \subseteq \mathbb{P}(R_{j+k})$ such that, for all $[h] \in U_3$, we have $(I_{X'})_{2j+2k} = (I_X + \langle h \rangle)_{2j+2k}$, which implies that $h_{X'}(2j+2k) = h_X(2j+2k) - h_X(j+k)$. The triple intersection $U_1 \cap U_2 \cap U_3$ is nonempty, so Proposition 3.2 (ii) yields the inequality $h_{X'}(k) + h_{X'}(j+k) \leq h_{X'}(2j+2k)$. By construction, we have $h_{X'}(j+k) \leq h_X(j+k) - 1$ and $h_{X'}(i) \leq h_X(i)$ for all $i < j+k$. Therefore, we conclude that $h_X(k) + 2h_X(j+k) - 1 \leq h_X(2j+2k)$ when the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. \square

The hypothesis in Theorem 4.3 may be recast using alternative numerical invariants. With this in mind, set $e_i(X) := \max\{i \in \mathbb{Z} : H^i(\mathbb{P}^n, \mathcal{I}_X(i)) \neq 0\}$, so that the Castelnuovo–Mumford regularity of \mathcal{I}_X equals $\max\{e_i(X) + i : i \in \mathbb{Z}\}$; compare with Theorem 4.3 in [E].

Corollary 4.4. *Fix a positive integer j and a nonnegative integer k . Let $X \subseteq \mathbb{P}^n$ be a totally-real curve of degree d and arithmetic genus p_a , and assume that*

$$k \geq \max\left\{e_1(X) + 1, \frac{1}{2}e_2(X) + \frac{1}{2} - j, \frac{2p_a}{d}\right\}.$$

If $f \in P_{X,2j}$, then there exists a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$.

Proof. When $j+k \geq e_1(X) + 1$, we have $H^1(\mathbb{P}, \mathcal{I}_X(j+k)) = (M_X)_{j+k} = 0$; compare with Remark 4.2. The Hilbert polynomial of X equals $p_X(i) = di + (1 - p_a)$ and satisfies

$$h_X(i) - p_X(i) = \dim H^2(\mathbb{P}^2, \mathcal{I}_X(i)) - \dim H^1(\mathbb{P}^2, \mathcal{I}_X(i)),$$

so we see that $h_X(i) = p_X(i)$ for all $i > \max\{e_1(X), e_2(X)\}$ and $h_X(i) \geq p_X(i)$ for all $i > e_1(X)$. Hence, if $k \geq e_1(X) + 1$ and $2j+2k \geq e_2(X) + 1$, then the inequality $k \geq \frac{2p_a}{d}$ or $k > \frac{2p_a-1}{d}$ is equivalently to $p_X(2j+2k) < 2p_X(j+k) + p_X(k) - 1$ and $h_X(2j+2k) < 2h_X(j+k) + h_X(k) - 1$. Therefore, Theorem 4.3 establishes the corollary. \square

For a second version, set $r(X) := \min\{j \in \mathbb{Z} : h_X(i) = p_X(i) \text{ for all } i \geq j\}$ where $p_X(i)$ denotes the Hilbert polynomial of X . This numerical invariant is sometimes called the Hilbert regularity of X or the index of regularity for X . A curve $X \subset \mathbb{P}^n$ is nondegenerate if it is not contained in a hyperplane.

Corollary 4.5. *Fix a positive integer j and a nonnegative integer k . Let $X \subset \mathbb{P}^n$ be a nondegenerate totally-real curve of degree d and arithmetic genus p_a , and assume that $k \geq \max\{r(X), \frac{2p_a}{d}\}$. If $f \in P_{X,2j}$, then there is a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$.*

Proof. The inequalities $k \geq r(X)$ and $k \geq \frac{2p_a}{d}$ yield $h_X(j+k) = p_X(j+k) = d(j+k) + (1 - p_a)$ and $d(j+k) - p_a = dj + dk - p_a \geq dj + p_a \geq 2$ respectively. By hypothesis, the line bundle $\mathcal{O}_X(1)$ is very ample and $j+k \geq 1$, so the complete linear series $|\mathcal{O}_X(j+k)|$ defines a closed immersion $\varphi: X \rightarrow \mathbb{P}^{d(j+k)-p_a}$. If $Y := \varphi(X)$, then a generic hyperplane section Y' of the curve Y consists of $d(j+k)$ points, any $d(j+k) - p_a$ of which are linearly independent; see the General Position Theorem on page 109 in [AC+]. Employing the inequality $k \geq \frac{2p_a}{d}$ as second time, we observe that $d(j+k) < d(j+k) + (dk - 2p_a) + dj - 1 = 2(d(j+k) - p_a - 1) + 1$. Hence, the Lemma on page 115 in [AC+] establishes that the points in Y' impose independent conditions on homogeneous polynomials of degree 2, and Corollary 4.7 in [E] shows that $\mathcal{I}_{Y'}$ is 2-regular. In particular, we obtain $H^1(\mathbb{P}^{d(j+k)-p_a}, \mathcal{I}_{Y'}(1)) = 0$. Since X is nondegenerate, the curve Y is also nondegenerate and we also have $H^1(\mathbb{P}^{d(j+k)-p_a}, \mathcal{I}_Y) = 0$. The long exact sequence in cohomology arising from the short exact sequence $0 \rightarrow \mathcal{I}_Y(-1) \rightarrow \mathcal{I}_Y \rightarrow \mathcal{I}_{Y'} \rightarrow 0$ implies that $0 = H^1(\mathbb{P}^{d(j+k)-p_a}, \mathcal{I}_Y(1)) = H^1(\mathbb{P}^n, \mathcal{I}_X(j+k))$. Since $k \geq r(X)$, we have $h_X(i) = p_X(i)$ for all $i \geq k$ and the inequality $k \geq \frac{2p_a}{d}$ is equivalent to $h_X(2j+2k) < 2h_X(j+k) + h_X(k) - 1$, as in the proof of Corollary 4.4. Therefore, Theorem 4.3 establishes the corollary. \square

We illustrate these corollaries for two classic families of curves.

Example 4.6 (Complete intersection curves). Consider a totally-real complete intersection curve $X \subseteq \mathbb{P}^n$ cut out by forms of degree d_1, d_2, \dots, d_{n-1} for $1 \leq i \leq n-1$ where at least one d_i is greater than 1. This curve is arithmetically Cohen–Macaulay, so $e_1(X) = -\infty$; compare with Remark 4.2. By breaking the minimal free resolution of \mathcal{I}_X (which is a Koszul complex) into short exact sequences and knowing the cohomology of line bundles on projective space, we deduce

that $r(X) = e_2(X) = d_1 + d_2 + \cdots + d_{n-1} - n - 1$. As in Example 1.5.1 in [M], the degree of X is $d_1 d_2 \cdots d_{n-1}$ and the arithmetic genus is $\frac{1}{2}(d_1 d_2 \cdots d_{n-1})(d_1 + d_2 + \cdots + d_{n-1} - n - 1) + 1$. Assuming that $k \geq d_1 + d_2 + \cdots + d_{n-1} - n$, Corollary 4.4 or Corollary 4.5 establish that, for all $f \in P_{X,2j}$, there exists a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$. \diamond

Example 4.7 (Planar curves). If $X \subset \mathbb{P}^2$ is a planar curve of degree d at least 2 and $k \geq d - 2$, then Example 4.6 implies that, for all $f \in P_{X,2j}$, there is a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$. \diamond

Although Example 5.3 shows that this degree bound from Corollary 4.5 is sharp on some planar curves, the next example demonstrates that this is not always the case. Moreover, it illustrates how our techniques yield sharper bounds when the convex algebraic geometry of the underlying variety is well understood.

Example 4.8 (Non-optimality for planar curves). Let $X \subset \mathbb{P}^2$ be a rational quartic curve with a real parametrization and a real triple point. For instance, the curve X could be the image of the map $[x_0 : x_1] \mapsto [x_0^2 x_1(x_0 - x_1) : x_0 x_1^2(x_0 - x_1) : x_0^4 + x_1^4]$ where $[1 : 0], [0 : 1], [1 : 1] \in \mathbb{P}^1$ are all sent to $[0 : 0 : 1] \in \mathbb{P}^2$. We claim that, for all $f \in P_{X,2}$, there exists a nonzero $g \in \Sigma_{X,2}$ such that $fg \in \Sigma_{X,4}$.

We first reduce the claim to showing that a generic linear functional $\ell \in R_4^*$ can be written as conjugate-invariant linear combination of at most 8 point evaluations on $X(\mathbb{C})$. If the claim is false, then there exists a linear functional $\ell \in R_4^*$ that strictly separates $\Sigma_{X,4}$ and $f \cdot \Sigma_{X,2}$. We may assume that $\ell \in R_4^*$ is a generic linear functional, because $\Sigma_{X,4}$ and $f \cdot \Sigma_{X,2}$ are pointed cones. Since $h_X(2) = 6$ and $h_X(1) = 3$, the affine hulls of $\Sigma_{X,4}$ and $f \cdot \Sigma_{X,2}$ have dimension 6 and 3 respectively. As analysis of symmetric forms arising from point evaluations appearing in the proof of Proposition 3.2 indicates, the number of real point evaluations with positive coefficients plus the number of pairs of complex point evaluations is at least 6 and the number of real point evaluations with negative coefficients plus the number of pairs of complex point evaluations is at least 3. However, if $\ell \in R_4^*$ is a conjugate-invariant linear combination of at most 8 point evaluations, then we obtain a contradiction.

It remains to show that a generic linear functional $\ell \in R_4^*$ is a conjugate-invariant linear combination of at most 8 point evaluations on $X(\mathbb{C})$. The curve X is a projection of the rational normal quartic curve $\check{X} \subset \mathbb{P}^4$. It follows that there is a linear surjection $\rho : \check{R}_4^* \rightarrow R_4^*$ sending point evaluations on \check{X} to point evaluations on X . Hence, it suffices to prove that, for a generic $\ell \in R_4^*$, there exists a linear functional $\check{\ell} \in \rho^{-1}(\ell)$ that is a conjugate-invariant linear combination of at most 8 points evaluations on $\check{X}(\mathbb{C})$.

By construction, the \mathbb{R} -vector space \check{R}_4 is isomorphic to $\mathbb{R}[x_0, x_1]_{16}$. Thus, a generic linear functional $\check{\ell} \in \check{R}_4$ can be written as a conjugate-invariant linear combination of at most 9 point evaluations; see Lemma 1.33 in [IK]. Moreover, $\check{\ell} \in \check{R}_4$ can be written as a conjugate-invariant linear combination of 9 point evaluations if and only if the corresponding (8×8) -catelecticant matrix is invertible; see Theorem 1.44 or the second paragraph on page 28 in [IK]. A general element in \check{R}_4 , for which the corresponding (8×8) -catelecticant matrix is not invertible, is a conjugate-invariant linear combination of 8 point evaluations. Hence, it is enough to show that there exists $\check{\ell} \in \rho^{-1}(\ell)$ for which the corresponding catelecticant is not invertible. Since three points of \check{X} are

mapped to the same point in X , there exists a linear functional $\check{\ell}' \in \rho^{-1}(0)$ such that corresponding (8×8) -catalecticant matrix has rank 3; compare with Theorem 1.43 in [IK]. Choose an arbitrary linear functional $\check{\ell}'' \in \rho^{-1}(\ell)$ and consider the pencil $\check{\ell}'' + \lambda \check{\ell}'$ where $\lambda \in \mathbb{R}$. The determinant of (8×8) -catalecticant matrix corresponding to $\check{\ell}'' + \lambda \check{\ell}'$ is a polynomial of degree 3 in λ . Since every real polynomial of degree 3 has at least one real root, we conclude that there is a value for $\lambda \in \mathbb{R}$ such that the linear functional $\check{\ell} := \check{\ell}'' + \lambda \check{\ell}'$ is a conjugate-invariant linear combination of at most 8 points evaluations on $\check{X}(\mathbb{C})$. \diamond

For a nondegenerate curve, we also give a uniform bound depending only on the degree.

Corollary 4.9. *Fix a positive integer j and a nonnegative integer k . Let $X \subset \mathbb{P}^n$ be a nondegenerate totally-real curve of degree d , and assume that $k \geq d - n + 1$. If $f \in P_{X,2j}$, then there exists a nonzero $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$.*

Proof. Theorem 1.1 in [GLP] proves that the Castelnuovo–Mumford regularity of \mathcal{S}_X is at most $d - n + 2$, so we have $e_1(X) \leq d - n$ and $e_2(X) \leq d - n - 1$. Theorem 3.2 in [N] establishes that

$$p_a \leq \begin{cases} \binom{d-2}{2} - (n-3) & \text{if } d \geq 3 \\ 2 - n & \text{if } d = 2, \end{cases}$$

from which we conclude that $\frac{2p_a}{d} \leq d - n + 1$. Thus, the claim follows from Corollary 4.4. \square

When the Hilbert functions of iterated hypersurface sections can be controlled, the techniques used to prove Theorem 4.3 also apply to higher-dimensional varieties. If a homogeneous polynomial is strictly positive on a totally-real variety, then the associated hypersurface section has no real points. Focusing on non-totally-real projective varieties is unexpectedly the key insight needed to establish our higher-dimensional results.

Lemma 4.10. *Fix a positive integer j , let $X \subseteq \mathbb{P}^n$ be an m -dimensional variety that is not totally-real, and assume that X is arithmetically Cohen–Macaulay. If $h_X(2j) < \binom{m+2}{1} h_X(j) - \binom{m+2}{2}$, then the cone $\Sigma_{X,2j}$ contains a line.*

Proof. To obtain a contradiction, suppose that the cone $\Sigma_{X,2j}$ contains no lines. Hence, Remark 2.7 shows that $\Sigma_{X,2j}$ is pointed. We begin by proving that there exists $h_1, h_2, \dots, h_m \in S_j$ such that $Z := X \cap V(h_1, h_2, \dots, h_m)$ is a reduced set of non-real points with the uniform position property. To achieve this, observe that Theorem 3.1 implies that the set of homogeneous polynomials $h \in S_j$, such that $h \notin I_X$, $X' := X \cap V(h) \subset \mathbb{P}^n$, and the cone $\Sigma_{X',2j}$ is pointed, contains a nonempty Euclidean open subset $U_1 \subseteq \mathbb{P}(R_j)$. Next, Bertini’s Theorem (see Théorème 6.3 in [J]) establishes that a general hypersurface section of a geometrically integral variety of dimension at least 2 is geometrically integral and that a general hypersurface section of a geometrically reduced variety is geometrically reduced. Thirdly, the hypothesis that X is arithmetically Cohen–Macaulay implies that X' is also arithmetically Cohen–Macaulay and $h_{X'}(k) = h_X(k) - h_X(k - j)$ for all $k \in \mathbb{Z}$. Finally, a general hypersurface section of non-totally-real variety is also not totally-real, and a general hypersurface section of a non-totally-real curve consists of non-real points. Combining these

four observations, we deduce that there exists homogeneous polynomials $h_1, h_2, \dots, h_m \in S_j$ such that the intersection $Z := X \cap V(h_1, h_2, \dots, h_{m-1})$ has the desired properties. As the cone $\Sigma_{Z,2j}$ is pointed, Proposition 3.2 (i) now shows that $h_Z(j) \leq \lceil \frac{1}{2} h_Z(2j) \rceil$ which yields $2h_Z(j) \leq h_Z(2j) + 1$. Since we have both $h_Z(j) = h_X(j) - m$ and $h_Z(2j) = h_X(2j) - m h_X(j) + \binom{m}{2}$, it follows that $\binom{m+2}{1} h_X(j) - \binom{m+2}{2} \leq h_X(2j)$ which gives the required contradiction. \square

The inequality in Lemma 4.10 has an elegant restatement in terms of the Artinian reduction of X .

Remark 4.11. If $h_{Z'}: \mathbb{Z} \rightarrow \mathbb{Z}$ is the Hilbert function of the Artinian quotient of R by a maximal regular sequence of degree j , then we have $h_{Z'}(k) = h_Z(k) - h_Z(k-j)$ where Z is the arithmetically Cohen–Macaulay variety defined in the antepenultimate sentence of the proof of Lemma 4.10. Hence, the inequality $h_X(2j) < \binom{m+2}{1} h_X(j) - \binom{m+2}{2}$ is equivalent to the inequality $h_{Z'}(2j) < h_{Z'}(j)$.

In a special case, the inequality in Lemma 4.10 may also be expressed in terms other invariants.

Remark 4.12. If $X \subseteq \mathbb{P}^n$ is nondegenerate, then we have $h_X(1) = n + 1$. Lemma 3.1 in [BSV] establishes that the quadratic deficiency $\varepsilon(X)$ equals $h_X(2) - (m+1)(n+1) + \binom{m+1}{2}$. Hence, the Addition Formula for binomial coefficients gives

$$\begin{aligned} h_X(2) - \binom{m+2}{1} h_X(1) + \binom{m+2}{2} &= [h_X(2) - \binom{m+1}{1} h_X(1) + \binom{m+1}{2}] - [\binom{m+1}{0} h_X(1) - \binom{m+1}{1}] \\ &= \varepsilon(X) - \text{codim}(X), \end{aligned}$$

so the inequality in Lemma 4.10 becomes $\varepsilon(X) < \text{codim}(X)$ when X is nondegenerate and $j = 1$.

Lemma 4.10 shows that there exists a nontrivial sum of squares equal to zero. Exploiting this observation, we can prove a higher-dimensional analogue of Theorem 4.3.

Theorem 4.13. Fix a positive integer j and a nonnegative integer k . Let $X \subseteq \mathbb{P}^n$ be a totally-real variety with dimension m . Assume that X is arithmetically Cohen–Macaulay and that

$$h_X(2j+2k) < \binom{m+1}{1} (h_X(j+k) - h_X(k-j)) + h_X(2k) - \binom{m+1}{2}.$$

If $f \in P_{X,2j}$, then there exists a nonzero $g \in R_{2k}$ such that $fg \in \Sigma_{X,2j+2k}$.

Proof. With the aim of producing a contradiction, suppose that, for all nonzero $g \in R_{2k}$, we have $fg \notin \Sigma_{X,2j+2k}$. This means that the linear subspace $f \cdot R_{2k} := \{fg \in R_{2j+2k} : g \in R_{2k}\} \subset R_{2j+2k}$ intersects the cone $\Sigma_{X,2j+2k}$ only at the origin. As X is totally-real, Corollary 2.8 establishes that the cone $\Sigma_{X,2j+2k}$ is pointed and, in particular, closed. Hence, there exists a Euclidean open neighbourhood U of $f \in R_{2j}$ such that, for all $h \in U$ and all nonzero $g \in R_{2k}$, we have $hg \notin \Sigma_{X,2j+2k}$. Bertini’s Theorem (see Théorème 6.3 in [J]) establishes that a general hypersurface section of a geometrically reduced variety is geometrically reduced. The cone $P_{X,2j}$ is full-dimensional, so there is a general $h \in U \cap P_{X,2j}$ such that $X' := X \cap V(h)$ and $R' = R/\langle h \rangle = S/(I_X + \langle h \rangle)$. Every real zero of h must be contained in the singular locus of X' , because $h \in P_{X,2j}$. As X' is reduced, its singular locus is a proper Zariski closed subset, which implies that X' is not totally real. Since X is arithmetically Cohen–Macaulay, the variety X' is also arithmetically Cohen–Macaulay and $h_{X'}(i) = h_X(i) - h_X(i-2j)$. From $h_X(2j+2k) < \binom{m+1}{1} (h_X(j+k) - h_X(k-j)) + h_X(2k) - \binom{m+1}{2}$,

we obtain $h_{X'}(2j+2k) < \binom{(m-1)+2}{1} h_{X'}(j+k) - \binom{(m-1)+2}{2}$. Hence, Lemma 4.10 shows that the cone $\Sigma_{X',2j+2k}$ contains a line. Applying Proposition 2.5, there exists nonzero $g'_1, g'_2, \dots, g'_s \in R'_{j+k}$ such that $(g'_1)^2 + (g'_2)^2 + \dots + (g'_s)^2 = 0$. Lifting this equation to the ring R , we see that there are $g_1, g_2, \dots, g_s \in R_{j+k}$ such that $g_1^2 + g_2^2 + \dots + g_s^2 \in \langle h \rangle$. However, this contradicts the fact that $h \in U$. Therefore, we conclude that there exists a nonzero $g \in R_{2k}$ such that $fg \in \Sigma_{X,2j+2k}$. \square

Remark 4.14. Suppose that $f \in P_{X,2j}$ is strictly positive on $X(\mathbb{R})$ or, more generally, that the subset $X(\mathbb{R}) \setminus V(f)$ is dense in the Euclidean topology. For instance, the second condition automatically holds when f is nonzero and $X(\mathbb{R})$ a cone over a manifold in which all of the connected components have the same dimension. With this extra hypothesis, the nonzero $g \in R_{2k}$ described in Theorem 4.13 must be nonnegative.

Remark 4.15. Suppose that $f \in P_{X,2j}$ is strictly positive on $X(\mathbb{R})$. If one allows the degree of the nonzero multiplier $g \in R_{2k}$ to be greater than or equal to the degree of $f \in P_{X,2j}$, then one obtains a frivolous sum-of-squares representation $fg = f^2h \in \Sigma_{X,2j+2k}$ by choosing $g := fh$ where $h \in \Sigma_{X,2k-2j}$. However, the products $fg \in \Sigma_{X,2j+2k}$ arising from Theorem 4.13 are never of this frivolous form, because Lemma 4.10 shows that they are lifted from a nontrivial sum-of-squares modulo f .

The next four examples showcase the most interesting applications of Theorem 4.13. In these examples, we also obtain simple explicit degree bounds on the sum-of-squares multipliers.

Example 4.16 (Nonnegative quadratic forms on varieties of minimal degree). Fix $j = 1$ and $k = 0$. Let $X \subseteq \mathbb{P}^n$ be a totally-real variety of minimal degree. In other words, the variety X is nondegenerate and $\deg(X) = 1 + \text{codim}(X) = 1 + n - m$ where $m := \dim(X)$. The classification of varieties of minimal degree (see Theorem 1 in [EH]) implies that X is arithmetically Cohen–Macaulay and $\sum_{i \in \mathbb{Z}} h_X(i) t^i = (1 + (n-m)t)(1-t)^{-(m+1)}$. Hence, the Generalized Binomial Theorem establishes that $h_X(i) = \binom{m+i}{m} + (n-m)\binom{m+i-1}{m}$ for $i \geq 1-m$. It follows that

$$\begin{aligned} & \binom{m+1}{1} (h_X(j+k) - h_X(k-j)) + h_X(2k) - \binom{m+1}{2} - h_X(2j+2k) \\ &= \binom{m+1}{1} \left(\binom{m+1}{m} + (n-m)\binom{m}{m} \right) + 1 - \binom{m+1}{2} - \left(\binom{m+2}{m} + (n-m)\binom{m+1}{m} \right) = 1 > 0, \end{aligned}$$

so Theorem 4.13 shows that $P_{X,2} = \Sigma_{X,2}$. This gives another proof of Proposition 4.1 in [BSV]. \diamond

Example 4.17 (Nonnegative forms on surfaces of minimal degree). Fix $j \geq 1$ and $k = j-1$. Let $X \subseteq \mathbb{P}^n$ be a totally-real surface of minimal degree. As in Example 4.16, X is arithmetically Cohen–Macaulay, and we have $h_X(i) = \binom{i+2}{2} + (n-2)\binom{i+1}{2}$ for $i \geq -1$. Since

$$\begin{aligned} & \binom{2+1}{1} (h_X(j+k) - h_X(k-j)) + h_X(2k) - \binom{2+1}{2} - h_X(2j+2k) \\ &= 3(h_X(2j-1) - h_X(-1)) + h_X(2j-2) - 3 - h_X(4j-2) = 4j-3 > 0, \end{aligned}$$

Theorem 4.13 shows that, for all $f \in P_{X,2j}$, there exists a nonzero $g \in R_{2j-2}$ such that $fg \in \Sigma_{X,4j-2}$. Remark 4.14 also implies that $g \in P_{X,2j-2}$. Because Example 4.16 proves that we have $g \in \Sigma_{X,2j-2}$

when $j = 1$, an induction on j establishes that, for all $f \in P_{X,2j}$, there exists a nonzero $h \in \Sigma_{X,j^2-j}$ such that $fh \in \Sigma_{X,j^2+j}$. \diamond

Example 4.18 (Nonnegative forms on the projective plane). Fix $j \geq 2$ and $k = j - 2$. The variety \mathbb{P}^2 is arithmetically Cohen–Macaulay and $h_{\mathbb{P}^2}(i) = \binom{i+2}{2}$ for $i \geq -2$. It follows that

$$\begin{aligned} & \binom{2+1}{1} (h_{\mathbb{P}^2}(j+k) - h_{\mathbb{P}^2}(k-j)) + h_{\mathbb{P}^2}(2k) - \binom{2+1}{2} - h_{\mathbb{P}^2}(2j+2k) \\ &= 3(h_{\mathbb{P}^2}(2j-2) - h_{\mathbb{P}^2}(-2)) + h_{\mathbb{P}^2}(2j-4) - 3 - h_{\mathbb{P}^2}(4j-4) = 2j-3 > 0, \end{aligned}$$

so Theorem 4.13 and Remark 4.14 show that, for all $f \in P_{\mathbb{P}^2,2j}$, there exists a nonzero $g \in P_{\mathbb{P}^2,2j-4}$ such that $fg \in \Sigma_{\mathbb{P}^2,4j-4}$. In particular, this re-establishes a result of Hilbert (see [Hil] or Theorem 2.6 in [BI]). As in Example 4.17, an induction on j proves that

- for all $f \in P_{\mathbb{P}^2,4j}$, there exists a nonzero $h \in \Sigma_{\mathbb{P}^2,2j^2-2j}$ such that $fh \in \Sigma_{\mathbb{P}^2,2j^2+2j}$, and
- for all $f \in P_{\mathbb{P}^2,4j-2}$, there exists a nonzero $h \in \Sigma_{\mathbb{P}^2,2j^2-4j+2}$ such that $fh \in \Sigma_{\mathbb{P}^2,2j^2}$.

Since $P_{\mathbb{P}^2,6} \neq \Sigma_{\mathbb{P}^2,6}$, this degree bound is sharp for $f \in P_{\mathbb{P}^2,6}$ and Example 5.17 shows that it is also sharp for $f \in P_{\mathbb{P}^2,8}$. \diamond

Example 4.19 (Nonnegative forms on some surfaces of almost minimal degree). Fix $j \geq 1$ and $k = j$. Let $X \subset \mathbb{P}^n$ be a totally-real surface that is arithmetically Cohen–Macaulay and satisfies $\sum_{i \in \mathbb{Z}} h_X(i)t^i = (1 + c_1t + c_2t^2 + c_3t^3 + c_4t^4)(1-t)^{-3}$ for some $c_1, c_2, c_3, c_4 \in \mathbb{Q}$. The Generalized Binomial Theorem yields $h_X(i) = \binom{i+2}{2} + c_1 \binom{i+1}{2} + c_2 \binom{i}{2} + c_3 \binom{i-1}{2} + c_4 \binom{i-2}{2}$ for all $i \geq 2$, so it follows that

$$\begin{aligned} & \binom{2+1}{1} (h_X(j+k) - h_X(k-j)) + h_X(2k) - \binom{2+1}{2} - h_X(2j+2k) \\ &= 3(h_X(2j) - h_X(0)) + h_X(2j) - 3 - h_X(4j) = 2(c_1 - c_2 - 3c_3 - 5c_4 + 3)j + 3(c_3 + 3c_4 - 1). \end{aligned}$$

Thus, if $2c_1 + 3 > 2c_2 + 3c_3 + c_4$, then Theorem 4.13 shows that, for all $f \in P_{X,2j}$, there exists a nonzero $g \in R_{2j}$ such that $fg \in \Sigma_{X,4j}$. For instance, if X is a totally-real surface of almost minimal degree that is arithmetically Cohen–Macaulay (in other words, the surface X is nondegenerate, arithmetically Cohen–Macaulay, and $\deg(X) = 2 + \text{codim}(X) = n$), then we have $c_1 = n - 2$, $c_2 = 1$, $c_3 = 0$, and $c_4 = 0$, which implies that $2c_1 + 3 = 2n - 1 > 2 = 2c_2 + 3c_3 + c_4$. By Remark 4.15, this certificate is not frivolous. \diamond

5. LOWER BOUNDS FOR SUM-OF-SQUARES MULTIPLIERS

This final section establishes lower bounds on the minimal degree of a sum-of-squares multiplier. These degree bounds for the non-existence of sum-of-square multipliers prove the second half of our main theorems. For Harnack curves on smooth toric surfaces, these degree bounds for the existence of strict-separators are a perfect complement to our degree bounds for the existence of sum-of-square multipliers.

Our first lemma relates the zeros of a nonnegative element to the zeros of any sum-of-squares multiplier. For a closed point $p \in X$, let $d_p: R \rightarrow T_p^*(X)$ denote the derivation that sends $f \in R$ to the class of $f - f(p)$ in the Zariski cotangent space at p .

Lemma 5.1. *Fix a positive integer j and a nonnegative integer k . Let $X \subseteq \mathbb{P}^n$ be a totally-real projective variety, and consider $f \in P_{X,2j}$ and $g \in \Sigma_{X,2k}$ such that $fg \in \Sigma_{X,2j+2k}$. If the real point $p \in X(\mathbb{R})$ satisfies $f(p) = 0$ and $d_p(f) \neq 0$, then we have $g(p) = 0$ and $d_p(g) = 0$.*

Proof. Suppose $fg = h$ where $h := h_0^2 + h_1^2 + \cdots + h_s^2$ for $h_0, h_1, \dots, h_s \in R_{j+k}$. Since $f(p) = 0$, we see that $h(p) = 0$ and $h_j(p) = 0$ for all $0 \leq j \leq s$. Hence, the Leibniz Rule establishes that $0 = 2h_0(p)d_p(h_0) + 2h_1(p)d_p(h_1) + \cdots + 2h_s(p)d_p(h_s) = d_p(h) = f(p)d_p(g) + g(p)d_p(f)$. By hypothesis, we have $f(p) = 0$ and $d_p(f) \neq 0$, which implies that $g(p) = 0$. Since g is a sum of squares, we conclude that $d_p(g) = 0$, as we just did for h . \square

Remark 5.2. When $f \in P_{X,2j}$, the hypothesis that $d_p(f) \neq 0$ can only be satisfied if p is a singular point on the variety X .

Equipped with this lemma, we show that there exists a planar curve for which the bound on degree of multipliers given in Example 4.7 is tight.

Example 5.3 (Optimality for a planar curve). Let $X \subset \mathbb{P}^2$ be the rational tricuspidal quartic curve defined by the equation $(x_0^2 + x_1^2)^2 + 2x_2^2(x_0^2 + x_1^2) - \frac{1}{3}x_2^4 - \frac{8}{3}x_2(x_0^3 - 3x_0x_1^2) = 0$. This curve is called the deltoid curve and is parametrized by $t \mapsto [\frac{1}{3}(2\cos(t) + \cos(2t)) : \frac{1}{3}(2\sin(t) - \sin(2t)) : 1]$ in the affine plane $x_2 = 1$. In other words, the real points of X consist of the hypocycloid generated by the trace of a fixed point on a circle that rolls inside a larger circle with one-and-a-half times its radius. The three cusps occur at the points $[1 : 0 : 1]$, $[-\frac{1}{2} : \frac{\sqrt{3}}{2} : 1]$, $[-\frac{1}{2} : -\frac{\sqrt{3}}{2} : 1]$, corresponding to $t = 0, \frac{2\pi}{3}, \frac{4\pi}{3}$ respectively, and lie on the conic $x_2^2 - x_1^2 - x_0^2$.

Consider $f \in P_{X,2j}$ such that $d_p(f) \neq 0$ at each cusp p in X . For instance, the polynomial $(x_2^2 - x_1^2 - x_0^2)(x_0^2 + x_1^2 + x_2^2)^{j-1}$ is nonnegative on X and has nonzero derivations at each cusp p on X . Suppose that there exists a nonzero $g \in \Sigma_{X,2}$ such that $fg \in \Sigma_{X,2j+2}$. Lemma 5.1 implies that $g(p) = 0$ and $d_p(g) = 0$ at each cusp p of X . Expressing g as a sum of linear forms, it follows that each of these linear forms vanishes at all three cusps. Since the three cusps are not collinear, this is impossible. Therefore, for all nonzero $g \in \Sigma_{X,2}$, we conclude that $fg \notin \Sigma_{X,2j+2}$. \diamond

We next examine rational curves on a projective surface. A **surface** $Y \subseteq \mathbb{P}^n$ is a two-dimensional projective variety; for more information on algebraic surfaces, see [Be].

Lemma 5.4. *Let $Y \subseteq \mathbb{P}^n$ be a real surface and let X be a curve on Y . If X has j isolated real points p_1, p_2, \dots, p_j , then there exists $f \in P_{X,2j}$ such that $f(p_i) = 0$ and $d_{p_i}(f) \neq 0$ for all $1 \leq i \leq j$.*

Proof. Fix coordinates on \mathbb{P}^n such that the hyperplane $V(x_0)$ does not contain any isolated real points on X . For each isolated singular point $p_i \in X(\mathbb{R})$ where $1 \leq i \leq j$, let $\tilde{p}_i \in \mathbb{A}^{n+1}(\mathbb{R})$ be the affine representative in which the 0-th component equals 1. Choose a real point \tilde{q}_i in $V(x_0 - 1) \subset \mathbb{A}^{n+1}(\mathbb{R})$ such that the closed ball centered at \tilde{q}_i with radius $\varepsilon_i := \|\tilde{p}_i - \tilde{q}_i\|^2 > 0$ does not contain an affine representative \tilde{p} where $p \in X(\mathbb{R})$ except for the point \tilde{p}_i corresponding to an isolated real point. For $1 \leq i \leq j$, consider $\tilde{h}_i := (x_1 - (\tilde{q}_i)_1 x_0)^2 + (x_2 - (\tilde{q}_i)_2 x_0)^2 + \cdots + (x_n - (\tilde{q}_i)_n x_0)^2 - \varepsilon_i x_0^2 \in S_2$. If \tilde{h}_i maps to $h_i \in R_2$ under the canonical quotient map from S to R , then we have $h_i \in P_{X,2}$, $h_i(p_i) = 0$,

and $d_{p_i}(h_i) \neq 0$ by construction. Hence, the product $f := h_1 h_2 \cdots h_j \in R_{2j}$ satisfies the conditions in the first part of the lemma. \square

To obtain the desired bounds, we make additional assumptions on the surface and the curve. On a curve, an ordinary double point (also known as a node or an A_1 -singularity) is a point where a curve intersects itself so that the two branches of the curve have distinct tangent lines. There are two types of ordinary real double points: a **crossing** has two real branches and a **solitary point** has two imaginary branches that conjugate to each other. Hence, an isolated ordinary real double point is a solitary point. The following proposition is the basic source of our bounds for strict-separators.

Proposition 5.5. *Let $Y \subseteq \mathbb{P}^n$ be a real smooth rational surface such that the anti-canonical divisor is effective, and let H be a hyperplane section of Y . For some positive integer j , assume that there exists a section in $H^0(Y, \mathcal{O}_Y(jH))$ that defines a real rational curve $X \subset Y$ of degree d and arithmetic genus p_a . If X has p_a solitary points p_1, p_2, \dots, p_{p_a} , then there exists $f \in P_{X, 2j+2}$ such that $f(p_i) = 0$ and $d_{p_i}(f) \neq 0$ for all $1 \leq i \leq p_a$. Moreover, if the nonzero element $g \in \Sigma_{X, 2k}$ satisfies $fg \in \Sigma_{X, 2j+2k+2}$, then we have $k \geq \frac{2p_a}{d}$.*

Proof. Let K be the canonical divisor on Y . Since X is projective and the divisor jH is effective, Serre Duality (see Theorem I.11 in [Be]) shows that $H^2(Y, \mathcal{O}_Y(jH + K)) = H^0(Y, \mathcal{O}_Y(-jH)) = 0$. As Y is rational and the irregularity and geometric genus of a surface are birational invariants (see Proposition III.20 in [Be]), we have $H^1(Y, \mathcal{O}_Y) = 0$ and $H^2(Y, \mathcal{O}_Y) = H^0(Y, \mathcal{O}_Y(K)) = 0$, so the Euler–Poincaré characteristic $\chi(\mathcal{O}_Y)$ equals 1. Applying the Riemann–Roch Theorem (see Theorems I.12 and I.15 in [Be]), it follows that

$$\chi(\mathcal{O}_Y(jH + K)) = \chi(\mathcal{O}_Y) + \frac{1}{2}((jH + K)^2 - (jH + K) \cdot K) = 1 + \frac{1}{2}((jH)^2 + (jH) \cdot K) = p_a,$$

and we deduce that $\dim H^0(Y, \mathcal{O}_Y(jH + K)) \geq p_a$.

We first prove that the solitary points impose independent conditions by verifying that there is no nonzero section of $\mathcal{O}_Y(jH + K)$ which vanishes at any $p_a - 1$ solitary points of X and at any additional point $q \in X$. Suppose there exists a nonzero section of $\mathcal{O}_Y(jH + K)$ which vanishes at $p_a - 1$ solitary points of X and an additional point $q \in X$. Let \tilde{Y} be the blowing up of the surface Y at $p_a - 1$ solitary points and the point q ; the corresponding exceptional divisors are $E_1, E_2, \dots, E_{p_a-1}, F$. If this hypothetical section vanishes at the chosen $p_a - 1$ nodes of X and the point $q \in X$ with multiplicities m_i and r respectively, then the line bundle

$$\mathcal{O}_{\tilde{Y}}(jH + K - m_1 E_1 - m_2 E_2 - \cdots - m_{p_a-1} E_{p_a-1} - rF)$$

restricted to the proper transform of X in \tilde{Y} would also have a section. However, the degree of the restriction (see Lemma I.6 in [Be]) equals

$$\begin{aligned} (jH + K - m_1 E_1 - m_2 E_2 - \cdots - m_{p_a-1} E_{p_a-1} - rF) \cdot (jH - 2E_1 - 2E_2 - \cdots - 2E_{p_a-1} - F) \\ = 2(p_a - 1) - 2(m_1 + m_2 + \cdots + m_{p_a-1}) - r < 0, \end{aligned}$$

which yields the required contradiction.

To prove the first part, choose a nonzero section $f_1 \in H^0(Y, \mathcal{O}_Y(jH + K))$ that vanishes at the solitary points p_2, p_3, \dots, p_{p_a} . The previous paragraph ensures that $f_1(p_1) \neq 0$. Because the solitary points p_1, p_2, \dots, p_{p_a} are isolated and imposed independent conditions, there exists a section $f_2 \in H^0(Y, \mathcal{O}_Y(jH + K))$, a small perturbation of f_1 that does not vanish at any point in $X(\mathbb{R})$. Since $-K$ is effective, we may also choose a nonzero section $f_3 \in H^0(Y, \mathcal{O}_Y(-K))$. By construction, the section $f_1 f_2 f_3^2 \in H^0(Y, \mathcal{O}_Y(2jH))$ is greater than or equal to zero at all points in $X(\mathbb{R}) \setminus \{p_1\}$; see Section 5 in [BSV] for more information about the sign of a global sections. Applying Lemma 5.4, there exists $f_4 \in H^0(Y, \mathcal{O}_Y(2H))$ such that $f_4(p_1) = 0$ and $d_{p_1}(f_4) \neq 0$. Hence, the section $f := f_1 f_2 f_3^2 f_4 \in H^0(Y, \mathcal{O}_Y(2(j+1)H))$, which is the restriction to Y of a hypersurface of degree $2j+2$ in \mathbb{P}^n , is nonnegative on X and satisfies $f(p_i) = 0$ and $d_{p_i}(f) \neq 0$ for all $1 \leq i \leq p_a$.

For the second part, consider a nonzero element $g \in \Sigma_{X, 2k}$ such that $fg \in \Sigma_{X, 2j+2k}$. Lemma 5.1 establishes that $g(p_i) = 0$ and $d_{p_i}(g) = 0$ for $1 \leq i \leq j$. Fix an element \tilde{g} of degree $2k$ in the \mathbb{Z} -graded coordinate ring of Y that maps to $g \in R_{2k}$ under the canonical quotient homomorphism and consider the curve $Z \subset Y$ defined by \tilde{g} . Since the element g is nonzero in R_{2k} , the curve Z does not contain the curve X . Let \hat{Y} be the blowing up of the surface Y at the p_a solitary points p_1, p_2, \dots, p_{p_a} and let E_1, E_2, \dots, E_{p_a} be the corresponding exceptional divisors in \hat{Y} . The proper transforms $\hat{X} \subset \hat{Y}$ and $\hat{Z} \subset \hat{Y}$ of the curves $X \subset Y$ and $Z \subset Y$ are linearly equivalent to the divisor classes $D_{\hat{X}} := eH - 2E_1 - 2E_2 - \dots - 2E_{p_a}$ and $D_{\hat{Z}} := 2kH - m_1E_1 - m_2E_2 - \dots - m_{p_a}E_{p_a}$ for some $m_i \geq 2$. Since \hat{X} is irreducible, the degree of the line bundle $\mathcal{O}_{\hat{X}}(D_Y)$ is nonnegative. Hence, we obtain $0 \leq D_{\hat{Z}} \cdot D_{\hat{X}} = 2ekH^2 - 2(m_1 + m_2 + \dots + m_{p_a}) \leq 2kd - 4p_a$, which yields $k \geq \frac{2p_a}{d}$. \square

Remark 5.6. By modifying the third paragraph in the proof of Proposition 5.5, one can obtain slightly better bounds when the canonical divisor K is a multiple of the hyperplane section H . In particular, this applies for $Y = \mathbb{P}^2$.

Although Proposition 5.5 is the latent source for our sharpness results, it is technically difficult to apply because of its hypotheses. To address this challenge, we exhibit the appropriate rational curves on toric surfaces. To be more precise, consider a smooth convex lattice polygon $Q \subset \mathbb{R}^2$ and its associated nonsingular toric surface Y_Q . Fix a cyclic ordering for the edges of Q , let $u_1, u_2, \dots, u_m \in \mathbb{Z}^2$ be the corresponding primitive inner normal vectors to the edges, and let D_1, D_2, \dots, D_m be the corresponding irreducible torus-invariant divisors on Y_Q . The anti-canonical divisor on Y_Q is the effective divisor $D_1 + D_2 + \dots + D_m$. From the canonical presentation for the convex polytope $Q = \{v \in \mathbb{R}^2 : \langle v, u_i \rangle \geq -a_i \text{ for } 1 \leq i \leq m\}$, we obtain the very ample divisor $A_Q := a_1D_1 + a_2D_2 + \dots + a_mD_m$ on Y_Q . For more background on toric geometry, see Section 2.3 and Section 4.2 in [CLS].

As in Subsection 2.2 in [KO], we call the real connected components of a curve $X \subset Y_Q$ *ovals* and treat isolated real points as degenerate ovals. Following Definition 8 in [Br], a *Harnack curve* $X \subset Y_Q$ is the image of a real morphism $\xi : C \rightarrow Y_Q$ satisfying three conditions:

- (1) the smooth real curve C has the maximal number of ovals (namely, one more than the genus of the curve C);

- (2) there is a distinguished oval in $C(\mathbb{R})$ containing disjoint arcs $\Gamma_1, \Gamma_2, \dots, \Gamma_m$ such that, for all $1 \leq i \leq m$, we have $\xi^{-1}(D_i) \subseteq \Gamma_i$; and
- (3) the cyclic orientation on the arcs induced by the distinguished oval is exactly $[\Gamma_1, \Gamma_2, \dots, \Gamma_m]$.

These special curves are germane, because Theorem 10 in [Br] establishes that all of the singularities on a Harnack curve are solitary points. By modifying the technique in Subsection 4.1 of [KO] for \mathbb{P}^2 , we construct rational Harnack curves on smooth projective toric surfaces.

Proposition 5.7. *If $Q \subset \mathbb{R}^2$ is a smooth two-dimensional lattice polygon, then there exists a rational Harnack curve on the toric variety Y_Q which is linearly equivalent to the associated very ample divisor A_Q and has arithmetic genus equal to the number of interior lattice points in Q .*

Proof. Following [C], a map from \mathbb{P}^1 to the smooth toric variety Y_Q is determined by a collection of line bundles and sections on \mathbb{P}^1 that satisfy certain compatibility and non-degeneracy conditions. To describe the required map, fix disjoint arcs $\Gamma_1, \Gamma_2, \dots, \Gamma_m$ on the circle $\mathbb{P}^1(\mathbb{R})$ such that the induced cyclic orientation is $[\Gamma_1, \Gamma_2, \dots, \Gamma_m]$. The intersection product $e_i := A_Q \cdot D_i$, for each $1 \leq i \leq m$, equals the normalized lattice distance of the corresponding edge in the polytope Q . The Divergence Theorem shows that $e_1 \langle v, u_1 \rangle + e_2 \langle v, u_2 \rangle + \dots + e_m \langle v, u_m \rangle = 0$ for all $v \in \mathbb{Z}^2$, so the line bundles $\mathcal{O}_{\mathbb{P}^1}(e_1), \mathcal{O}_{\mathbb{P}^1}(e_2), \dots, \mathcal{O}_{\mathbb{P}^1}(e_m)$ satisfy the compatibility condition in Definition 1.1 in [C]. For all $1 \leq i \leq m$, choose distinct points $[c_{i,1} : 1], [c_{i,2} : 1], \dots, [c_{i,e_i} : 1] \in \Gamma_i$. Identifying global sections of $\mathcal{O}_{\mathbb{P}^1}(e_i)$ with homogeneous polynomials in $\mathbb{C}[x_0, x_2]_{e_i}$, we obtain the real sections $\prod_{j=1}^{e_i} (x_0 - c_{i,j} x_1) \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(e_i))$. Since we chose distinct points, no two sections vanish at the same point in \mathbb{P}^1 , so these sections satisfy the non-degeneracy condition in Definition 1.1 in [C]. Hence, Theorem 1.1 in [C] establishes that these line bundles and sections determine a real morphism $\xi : \mathbb{P}^1 \rightarrow Y_Q$ such that $\xi^{-1}(D_i) = \{[c_{i,1} : 1], [c_{i,2} : 1], \dots, [c_{i,e_i} : 1]\}$ for all $1 \leq i \leq m$. In other words, the image of ξ is a rational Harnack curve $X \subset Y_Q$. By construction, the curve X is also linearly equivalent to the divisor A_Q . Hence, Proposition 10.5.8 in [CLS] proves that the arithmetic genus of X equals the number of interior lattice points in Q . \square

Having assembled the necessary prerequisites, we now describe our lower bound on the degrees of sums-of-squares multipliers on curves.

Theorem 5.8. *For all $j \geq 2$, there exist smooth curves $X \subset \mathbb{P}^n$ and elements $f \in P_{X,2j}$ such that the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated for all $k < \frac{2p_a}{d}$ where d and p_a are the degree and genus of X respectively.*

Proof. Fix a smooth two-dimensional lattice polytope Q and let A_Q be the associated very ample divisor on the smooth toric variety Y_Q . Applying Proposition 5.7 to the dilated polytope $(j-1)Q$ gives a rational Harnack curve X on Y_Q of degree d defined by a section in $H^0(Y_Q, \mathcal{O}_{Y_Q}((j-1)A_Q))$. The number of singular points on X equals its arithmetic genus p_a and, as Theorem 10 in [Br] establishes, all of the singularities on X are solitary points. Hence, Proposition 5.5 shows that there exists an element $f \in P_{X,2j}$ such that, for all $k < \frac{2p_a}{d}$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. Asserting that the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated is an open condition in the Euclidean topology on the element $f \in R_{2j}$. Hence, we may assume that the given element f lies

in the interior of the cone $P_{X,2j}$. To finish the proof, we prove that, under small real perturbations of both X and f , the pertinent cones continue to be well-separated.

We first deform the singular Harnack curve X into a smooth Harnack curve X_ε . For brevity, let H denote the very ample divisor $(j-1)A_Q$. Fix a section $g_1 \in H^0(Y_Q, \mathcal{O}_{Y_Q}(H))$ defining X on Y_Q . Since H is very ample, we may choose a section $g_2 \in H^0(Y_Q, \mathcal{O}_{Y_Q}(H))$ that does not vanish at any solitary point of X , so the quotient g_1/g_2 is real-valued on $Y_Q \setminus V(g_2)$ and every solitary point of X is either a local maximum or local minimum. The product of sections defining the irreducible torus-invariant divisors determines a section $g_3 \in H^0(Y_Q, \mathcal{O}_{Y_Q}(-K))$, because the canonical divisor on the toric variety Y_Q is $K = -D_1 - D_2 - \cdots - D_m$. As the first paragraph in the proof of Proposition 5.5 establishes, the solitary points impose independent conditions on the sections of $\mathcal{O}_{Y_Q}(H+K)$. It follows that there exists a section $g_4 \in H^0(Y_Q, \mathcal{O}_{Y_Q}(H+K))$ such that the rational function g_3g_4/g_2 has prescribed values at the solitary points of X . In particular, we may choose the section g_4 so that g_3g_4/g_2 is negative at the local minima of the quotient g_1/g_2 and is positive at the local maxima of the quotient g_1/g_2 . For small enough $\varepsilon > 0$, we see that the section $g_1 + \varepsilon g_3g_4$ defines a smooth Harnack curve X_ε on Y_Q with arithmetic genus p_a . Moreover, the sections defining X_ε and X have the same degree, so we have $h_{X_\varepsilon}(i) = h_X(i)$ for all $i \in \mathbb{Z}$.

To deform the element $f \in P_{X,2j}$, choose a polynomial $\tilde{f} \in S_{2j}$ that maps to f under the canonical quotient homomorphism, set $e := h_X(2j+2k)$, and fix points p_1, p_2, \dots, p_e in X for which the linear functionals $p_1^*, p_2^*, \dots, p_e^*$, defined by point evaluation, form a basis for R_{2j+2k}^* . Since the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated, there exists a linear functional $\ell \in R_{2j+2k}^*$ satisfying $\ell(h) > 0$ for all nonzero $h \in \Sigma_{X,2j+2k}$ and $\ell(h) < 0$ for all nonzero $h \in f \cdot \Sigma_{X,2k}$. Hence, there are $\lambda_1, \lambda_2, \dots, \lambda_e \in \mathbb{R}$ such that $\ell = \lambda_1 p_1^* + \lambda_2 p_2^* + \cdots + \lambda_e p_e^*$. By choosing affine representatives $\tilde{p}_1, \tilde{p}_2, \dots, \tilde{p}_e \in \mathbb{A}^{n+1}$, we obtain $\tilde{\ell} := \lambda_1 \tilde{p}_1^* + \lambda_2 \tilde{p}_2^* + \cdots + \lambda_e \tilde{p}_e^*$ in S_{2j+2k}^* . There are two symmetric forms associated to the linear functional $\tilde{\ell}$: the first $\sigma_{j+k}^*(\tilde{\ell}): S_{j+k} \otimes_{\mathbb{R}} S_{j+k} \rightarrow \mathbb{R}$ is defined by $\tilde{h}_1 \otimes \tilde{h}_2 \mapsto \tilde{\ell}(\tilde{h}_1 \tilde{h}_2)$ and the second $\tau_j^*(\tilde{\ell}): S_k \otimes_{\mathbb{R}} S_k \rightarrow \mathbb{R}$ is defined by $\tilde{h}_1 \otimes \tilde{h}_2 \mapsto \tilde{\ell}(\tilde{f} \tilde{h}_1 \tilde{h}_2)$. The assertion that ℓ is a strict separator for the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ is equivalent to saying that symmetric form $\sigma_{j+k}^*(\tilde{\ell})$ is positive-semidefinite with $\text{Ker}(\sigma_{j+k}^*(\tilde{\ell})) = (I_X)_{j+k}$ and symmetric form $\tau_k^*(\tilde{\ell})$ is negative-semidefinite with $\text{Ker}(\tau_k^*(\tilde{\ell})) = (I_X)_k$. To build the applicable linear functional on the deformation X_ε , let q_1, q_2, \dots, q_e denote the points on X_ε corresponding to the fixed points p_1, p_2, \dots, p_e on X . Choose affine representatives $\tilde{q}_1, \tilde{q}_2, \dots, \tilde{q}_e \in \mathbb{A}^{n+1}$ and consider the linear functional

$$\tilde{\ell}_\varepsilon := \lambda_1 \tilde{q}_1^* + \lambda_2 \tilde{q}_2^* + \cdots + \lambda_e \tilde{q}_e^* \in S_{2j+2k}^*.$$

By construction, we have $(I_{X_\varepsilon})_{j+k} \subseteq \text{Ker}(\sigma_{j+k}^*(\tilde{\ell}_\varepsilon))$ and $(I_{X_\varepsilon})_k \subseteq \text{Ker}(\tau_k^*(\tilde{\ell}_\varepsilon))$. For sufficiently small $\varepsilon > 0$, the symmetric forms $\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)$ and $\tau_k^*(\tilde{\ell}_\varepsilon)$ are small perturbations of $\sigma_{j+k}^*(\tilde{\ell})$ and $\tau_k^*(\tilde{\ell})$ respectively. The rank of a symmetric form is lower semicontinuous, so we have both $\text{rank}(\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)) \geq \text{rank}(\sigma_{j+k}^*(\tilde{\ell}))$ and $\text{rank}(\tau_k^*(\tilde{\ell}_\varepsilon)) \geq \text{rank}(\tau_k^*(\tilde{\ell}))$. Because $h_{X_\varepsilon}(k) = h_X(k)$ and $h_{X_\varepsilon}(j+k) = h_X(j+k)$, it follows that $\text{Ker}(\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)) = (I_{X_\varepsilon})_{j+k}$ and $\text{Ker}(\tau_k^*(\tilde{\ell}_\varepsilon)) = (I_{X_\varepsilon})_k$. In addition, being positive-semidefinite or negative-semidefinite is an open condition in the Euclidean

topology, so the symmetric form $\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)$ is positive-semidefinite and symmetric form $\tau_k^*(\tilde{\ell}_\varepsilon)$ is negative-semidefinite. If f_ε denotes the image of \tilde{f} under the canonical quotient map from S to \mathbb{Z} -graded coordinate ring of X_ε , then we conclude that $\ell_\varepsilon := \lambda_1 q_1^* + \lambda_2 q_2^* + \cdots + \lambda_e q_e^*$ is a strict separator for the cones $\Sigma_{X_\varepsilon, 2j+2k}$ and $f_\varepsilon \cdot \Sigma_{X_\varepsilon, 2j+2k}$. \square

Remark 5.9. Although the smooth curves constructed in the proof of Theorem 5.8 have the maximal number of ovals, this is not necessary. By choosing the section g_4 so that $g_3 g_4 / g_2$ is positive at some local minima, or negative at some local maxima, of the quotient g_1 / g_3 , we can obtain smooth curves for which the number of ovals is anywhere between 1 and one more than the genus. In particular, Theorem 5.8 is remarkably insensitive to the topology of the real projective curve.

Remark 5.10. By applying the perturbation techniques from the proof of Theorem 5.8 to the tricuspidal curve in Example 5.3, we see that there are smooth planar curves for which the bound in Example 4.7 is tight.

For the smooth curves created in the proof of Theorem 5.8, both the degree and genus can be expressed as a functions of the parameter j . From these expressions, we see that, for all $j \geq 2$, there are smooth curves for which Theorem 5.8 is an exact counterpart to Corollary 4.5.

Example 5.11 (Curves with sharp bounds). Let $Q \subset \mathbb{R}^2$ be a smooth convex lattice polygon with an interior lattice point. Hence, we obtain a smooth toric variety $Y_Q \subset \mathbb{P}^n$ embedded by the very ample line bundle A_Q . The Ehrhart polynomial of Q equals the Hilbert polynomial of $Y_Q \subset \mathbb{P}^n$; see Proposition 9.4.3 in [CLS]. If $\text{area}(Q)$ denotes the standard Euclidean area of the polygon Q and $|\partial Q \cap \mathbb{Z}^2|$ counts the number of lattice points on its boundary ∂Q , then it follows that $p_{Y_Q}(i) = \text{area}(Q)i^2 + \frac{1}{2}|\partial Q \cap \mathbb{Z}^2|i + 1$; see Proposition 10.5.6 in [CLS].

Fix an integer j with $j \geq 2$. Since the smooth curves X appearing in the proof of Theorem 5.8 are defined by a section in $H^0(Y_Q, \mathcal{O}_{Y_Q}((j-1)A_Q))$, we have

$$p_X(i) = p_{Y_Q}(i) - p_{Y_Q}(i - (j-1)) = 2\text{area}(Q)(j-1)i + \frac{1}{2}|\partial Q \cap \mathbb{Z}^2|(j-1) - \text{area}(Q)(j-1)^2,$$

so the degree and genus of X are $2\text{area}(Q)(j-1)$ and $p_{Y_Q}(1-j)$ respectively. Amusingly, we have $\deg(X) = (j-1)\deg(Y_Q)$ and the genus equals the number of interior lattice points in the dilate $(j-1)Q$; see Theorem 9.4.2 in [CLS]. In addition, the equation for $p_X(i)$ implies that $r(X) = j-1-m$ where m is the largest nonnegative integer such that the dilate mQ does not contain any interior lattice points. Since a smooth polytope has at least three vertices, we have $3 \leq |\partial Q \cap \mathbb{Z}^2|$, $1 < \frac{1}{2}|\partial Q \cap \mathbb{Z}^2|(j-1)$, and

$$\begin{aligned} \left\lceil \frac{2p_a}{d} \right\rceil &= \left\lceil \frac{\text{area}(Q)(j-1)^2 - \frac{1}{2}|\partial Q \cap \mathbb{Z}^2|(j-1) + 1}{\text{area}(Q)(j-1)} \right\rceil \\ &\leq (j-1) + \left\lceil \frac{1 - \frac{1}{2}|\partial Q \cap \mathbb{Z}^2|(j-1)}{\text{area}(Q)(j-1)} \right\rceil \leq j-1. \end{aligned}$$

As Q has at least one interior lattice point, we also have $1 \leq p_{Y_Q}(-1) = \text{area}(Q) - \frac{1}{2}|\partial Q \cap \mathbb{Z}^2| + 1$, $1 < \text{area}(Q)$, and

$$\left\lfloor \frac{2p_a}{d} \right\rfloor \geq (j-1) - \left\lfloor \frac{\frac{1}{2}|\partial Q \cap \mathbb{Z}^2|}{\text{area}(Q)} \right\rfloor + \left\lfloor \frac{1}{\text{area}(Q)(j-1)} \right\rfloor = j-2.$$

Therefore, Theorem 5.8 proves that, for all $j \geq 2$, there exists smooth curves $X \subset \mathbb{P}^n$ and elements $f \in P_{X,2j}$ such that, for all $k < j-1$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. Conversely, Corollary 4.5 establishes that, for all $f \in P_{X,2j}$ and for all $k \geq j-1$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are not well-separated. \diamond

To be comprehensive, we also consider the smooth convex lattice polygons without an interior lattice point. From the classification of smooth toric surfaces (see Theorem 10.4.3 in [CLS]), we see that the polytopes omitted by Example 5.11 correspond to Hirzebruch surfaces and the projective plane. Using similar techniques to analyze these polytopes, we produce curves with sharp bounds contained in slightly smaller projective spaces.

Example 5.12 (Sharp bound for curves on Hirzebruch surfaces). For all $r, s \in \mathbb{N}$, consider the smooth lattice polygon $Q := \text{conv}\{(0,0), (s+1,0), (r+s+1,1), (0,1)\} \subset \mathbb{R}^2$. Since $|Q \cap \mathbb{Z}^2| = r+2s+4$, we obtain, for all $n \geq 3$, a Hirzebruch surface $Y_Q = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(r)) \subset \mathbb{P}^n$ embedded by the very ample line bundle A_Q . Fix an integer j with $j \geq 2$. Because we have $\text{area}(Q) = \frac{1}{2}r+s+1$ and $|\partial Q \cap \mathbb{Z}^2| = r+2s+4$, the calculations in Example 5.11 establish that, for the relevant curves $X \subset Y_Q$, we have

$$\frac{2p_a}{d} = \frac{(\frac{1}{2}r+s+1)(j-1)^2 - \frac{1}{2}(r+2s+4)(j-1) + 1}{(\frac{1}{2}r+s+1)(j-1)} = j-2 + \frac{2-j}{(\frac{1}{2}r+s+1)(j-1)},$$

and $j-3 < \lceil \frac{2p_a}{d} \rceil \leq j-2$. In addition, we have $r(X) = j-2$. Therefore, Theorem 5.8 proves that, for all $n \geq 3$ and for all $j \geq 2$, there exists smooth curves $X \subset \mathbb{P}^n$ and elements $f \in P_{X,2j}$ such that, for all $k < j-2$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. Conversely, Corollary 4.5 establishes that, for all $f \in P_{X,2j}$ and for all $k \geq j-2$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are not well-separated. \diamond

Example 5.13 (Sharp bounds for planar curves). Let $Q := \text{conv}\{(0,0), (1,0), (0,1)\} \subset \mathbb{R}^2$ be the standard simplex. Since $|Q \cap \mathbb{Z}^2| = 3$, we have $Y_Q = \mathbb{P}^2 \subseteq \mathbb{P}^2$ embedded by the very ample line bundle $A_Q = \mathcal{O}_{\mathbb{P}^2}(1)$. Fix an integer j with $j \geq 2$. Because we have $\text{area}(Q) = \frac{1}{2}$ and $|\partial Q \cap \mathbb{Z}^2| = 3$, the calculations in Example 5.11 establish that, for the relevant curves $X \subset Y_Q$, we have

$$\frac{2p_a}{d} = \frac{\frac{1}{2}(j-1)^2 - \frac{3}{2}(j-1) + 1}{\frac{1}{2}(j-1)} = j-4 + \frac{2}{j-1}.$$

When $j \geq 3$, we obtain $j-4 < \lceil \frac{2p_a}{d} \rceil \leq j-3$ and, when $j = 2$, we have $\frac{2p_a}{d} = 0$. In addition, we have $r(X) = j-3$. Therefore, Theorem 5.8 proves that, for all $j \geq 2$, there exists smooth curves $X \subset \mathbb{P}^2$ and elements $f \in P_{X,2j}$ such that, for all $0 \leq k < j-3$, the cones $\Sigma_{X,2j+2k}$ and

$f \cdot \Sigma_{X,2k}$ are well-separated. Conversely, Corollary 4.5 establishes that, for all $f \in P_{X,2j}$ and for all $k \geq \max\{j-3, 0\}$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are not well-separated. \diamond

Example 5.14 (Sharp bounds on the Veronese surface). Let $Q := \text{conv}\{(0,0), (2,0), (0,2)\} \subset \mathbb{R}^2$. Since $|Q \cap \mathbb{Z}^2| = 6$, we obtain the Veronese surface $Y_Q \subset \mathbb{P}^5$ embedded by the very ample line bundle $A_Q = \mathcal{O}_{\mathbb{P}^2}(2)$. Fix an integer j with $j \geq 2$. Because we have $\text{area}(Q) = 2$ and $|\partial Q \cap \mathbb{Z}^2| = 6$, the calculations in Example 5.11 establish that, for the relevant curves $X \subset Y_Q$, we have

$$\frac{2p_a}{d} = \frac{2(j-1)^2 - 3(j-1) + 1}{2(j-1)} = j - 2 + \frac{2-j}{2(j-1)},$$

and $j-3 < \lceil \frac{2p_a}{d} \rceil \leq j-2$. In addition, we have $r(X) = j-2$. Therefore, Theorem 5.8 proves that, for all $j \geq 2$, there exists smooth curves $X \subset \mathbb{P}^5$ and elements $f \in P_{X,2j}$ such that, for all $k < j-2$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. Conversely, Corollary 4.5 establishes that, for all $f \in P_{X,2j}$ and for all $k \geq j-2$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are not well-separated. \diamond

Proof of Theorem 1.1. Corollary 4.5 proves the first part. If one overlooks the parameter n , then Theorem 5.8 immediately proves the second part. By combining Example 5.12 and Example 5.13, it follows that the required curves and nonnegative elements exist for all $n \geq 2$. \square

We end this paper by lifting these degree bounds for strict-separators from curves to some surfaces. To accomplish this, we exploit the perturbation methods used in the proof of Theorem 5.8.

Proposition 5.15. *Fix a positive integer j and a nonnegative integer k . Let $X \subseteq \mathbb{P}^n$ be an arithmetically Cohen–Macaulay real projective variety and let X' be a hypersurface section of X of degree j . If there exists an element $f' \in P_{X',2j}$ such that the cones $\Sigma_{X',2j+2k}$ and $f' \cdot \Sigma_{X',2k}$ are well-separated, then there exists an element $f \in P_{X,2j}$ such that the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are also well-separated.*

Proof. We first lift f' to a nonnegative element on X . As observed in the proof of Proposition 5.8, asserting that the cones $\Sigma_{X',2j+2k}$ and $f' \cdot \Sigma_{X',2k}$ are well-separated is an open condition in the Euclidean topology on the element $f' \in R'_{2j}$. Hence, we may assume that f' is positive on $X'(\mathbb{R})$. Choose a homogeneous polynomial $\tilde{f}' \in S_{2j}$ that maps to f' under the canonical quotient homomorphism from S to $R' = S/I_{X'}$. By hypothesis, X' is a hypersurface section of X of degree j , so there is a nonzero polynomial $h \in S_j$ such that $X' = X \cap V(h) \subset \mathbb{P}^n$. Moreover, we have $I_{X'} = I_X + \langle h \rangle$, because X is arithmetically Cohen–Macaulay. Let $\tilde{X} \subseteq \mathbb{A}^{n+1}(\mathbb{R})$ be the affine cone of X and let $\mathbb{S}^n \subset \mathbb{A}^{n+1}(\mathbb{R})$ be the unit sphere. Since \tilde{f}' is positive on $X'(\mathbb{R})$, there exists a Euclidean neighbourhood U of $\mathbb{S}^n \cap V(h) \subset \mathbb{S}^n \cap \tilde{X}$ such that \tilde{f}' is positive on U . On the compact set $K := (\mathbb{S}^n \cap \tilde{X}) \setminus U$, the function h^2 is positive, so

$$\delta := \frac{\inf_K h^2}{\sup_K |\tilde{f}'|}$$

is positive real number. It follows that, for all $\lambda > \frac{1}{\delta}$, the polynomial $\tilde{f} := \tilde{f}' + \lambda h^2$ is positive on $X(\mathbb{R})$. Therefore, if f denotes the image of \tilde{f} under the canonical quotient homomorphism from S to $R = S/I_X$, then we deduce that $f \in P_{X,2j}$.

We next deform X' and f' . If $h = \sum_{|u|=2j} c_u x^u$ where $u \in \mathbb{N}^{n+1}$ and $c_u \in \mathbb{R}$, then consider the homogeneous polynomial $h_\varepsilon := \sum_{|u|=2j} (c_u + \varepsilon_u) x^u$ with $|\varepsilon_u| < \varepsilon$ created by perturbing the coefficients and the corresponding hypersurface section $X'_\varepsilon := X \cap V(h_\varepsilon) \subset \mathbb{P}^n$. Set $e := h_{X'}(2j+2k)$ and fix points p_1, p_2, \dots, p_e in X' for which the linear functionals $p_1^*, p_2^*, \dots, p_e^*$, defined by point evaluation, form a basis for $(R'_{2j+2k})^*$. Since the cones $\Sigma_{X', 2j+2k}$ and $f' \cdot \Sigma_{X', 2k}$ are well-separated, there exists a linear functional $\tilde{\ell} \in (R'_{2j+2k})^*$ satisfying $\tilde{\ell}(g) > 0$ for all nonzero $g \in \Sigma_{X', 2j+2k}$ and $\tilde{\ell}(g) < 0$ for all nonzero $g \in f' \cdot \Sigma_{X', 2k}$. It follows that there are real numbers $\lambda_1, \lambda_2, \dots, \lambda_e$ such that $\tilde{\ell} = \lambda_1 p_1^* + \lambda_2 p_2^* + \dots + \lambda_e p_e^*$. By choosing affine representatives $\tilde{p}_1, \tilde{p}_2, \dots, \tilde{p}_e \in \mathbb{A}^{n+1}$, we obtain $\tilde{\ell} := \lambda_1 \tilde{p}_1^* + \lambda_2 \tilde{p}_2^* + \dots + \lambda_e \tilde{p}_e^*$ in S_{2j+2k}^* . As in the proof of Theorem 5.8, there are two symmetric forms associated to the linear functional $\tilde{\ell}$: the first $\sigma_{j+k}^*(\tilde{\ell}) : S_{j+k} \otimes_{\mathbb{R}} S_{j+k} \rightarrow \mathbb{R}$ is defined by $\tilde{g}_1 \otimes \tilde{g}_2 \mapsto \tilde{\ell}(\tilde{g}_1 \tilde{g}_2)$ and the second $\tau_k^*(\tilde{\ell}) : S_k \otimes_{\mathbb{R}} S_k \rightarrow \mathbb{R}$ is defined by $\tilde{g}_1 \otimes \tilde{g}_2 \mapsto \tilde{\ell}(\tilde{f} \tilde{g}_1 \tilde{g}_2)$. The assertion that $\tilde{\ell}$ is a strict separator for the cones $\Sigma_{X', 2j+2k}$ and $f' \cdot \Sigma_{X', 2k}$ is equivalent to saying that symmetric form $\sigma_{j+k}^*(\tilde{\ell})$ is positive-semidefinite with $\text{Ker}(\sigma_{j+k}^*(\tilde{\ell})) = (I_{X'})_{j+k}$ and symmetric form $\tau_k^*(\tilde{\ell})$ is negative-semidefinite with $\text{Ker}(\tau_k^*(\tilde{\ell})) = (I_{X'})_k$. To build the applicable linear functional on a deformation X'_ε , let q_1, q_2, \dots, q_e denote the points on X'_ε corresponding to the fixed points p_1, p_2, \dots, p_e on X . Choose affine representatives $\tilde{q}_1, \tilde{q}_2, \dots, \tilde{q}_e \in \mathbb{A}^{n+1}$ and consider the linear functional $\tilde{\ell}_\varepsilon := \lambda_1 \tilde{q}_1^* + \lambda_2 \tilde{q}_2^* + \dots + \lambda_e \tilde{q}_e^*$ in S_{2j+2k}^* . By construction, we have $(I_{X_\varepsilon})_{j+k} \subseteq \text{Ker}(\sigma_{j+k}^*(\tilde{\ell}_\varepsilon))$ and $(I_{X_\varepsilon})_k \subseteq \text{Ker}(\tau_k^*(\tilde{\ell}_\varepsilon))$. For sufficiently small $\varepsilon > 0$, the symmetric forms $\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)$ and $\tau_k^*(\tilde{\ell}_\varepsilon)$ are small perturbations of $\sigma_{j+k}^*(\tilde{\ell})$ and $\tau_k^*(\tilde{\ell})$ respectively. The rank of a symmetric form is lower semicontinuous, so we have $\text{rank}(\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)) \geq \text{rank}(\sigma_{j+k}^*(\tilde{\ell}))$ and $\text{rank}(\tau_k^*(\tilde{\ell}_\varepsilon)) \geq \text{rank}(\tau_k^*(\tilde{\ell}))$. It follows that $\text{Ker}(\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)) = (I_{X'_\varepsilon})_{j+k}$ and $\text{Ker}(\tau_k^*(\tilde{\ell}_\varepsilon)) = (I_{X'_\varepsilon})_k$, because $h_{X'_\varepsilon}(k) = h_{X'}(k)$ and $h_{X'_\varepsilon}(j+k) = h_{X'}(j+k)$. In addition, being positive-semidefinite or negative-semidefinite is an open condition in the Euclidean topology, so the symmetric form $\sigma_{j+k}^*(\tilde{\ell}_\varepsilon)$ is positive-semidefinite and symmetric form $\tau_k^*(\tilde{\ell}_\varepsilon)$ is negative-semidefinite. If f'_ε denotes the image of \tilde{f} under the canonical quotient map from S to $R'_\varepsilon = S/(I_X + \langle h_\varepsilon \rangle)$, then we conclude that the linear functional $\ell_\varepsilon := \lambda_1 q_1^* + \lambda_2 q_2^* + \dots + \lambda_e q_e^*$ is a strict separator for the cones $\Sigma_{X'_\varepsilon, 2j+2k}$ and $f'_\varepsilon \cdot \Sigma_{X'_\varepsilon, 2k}$.

Lastly, suppose that there exists a nonzero $g \in \Sigma_{X, 2k}$ such that $fg \in \Sigma_{X, 2j+2k}$. By construction, the nonnegative element f restricts to f'_ε and the cones $\Sigma_{X'_\varepsilon, 2j+2k}$ and $f'_\varepsilon \cdot \Sigma_{X'_\varepsilon, 2k}$ are well-separated, so the multiplier g restricts to 0 on X'_ε . Equivalently, if g'_ε denotes the image of g under the canonical quotient map from R to $R'_\varepsilon = R/\langle h_\varepsilon \rangle$, then we have $g'_\varepsilon \in \langle h_\varepsilon \rangle$. Since this holds for all sufficiently small $\varepsilon \geq 0$, we see that $g = 0$ in R which is a contradiction. Therefore, the cones $\Sigma_{X, 2j+2k}$ and $f \cdot \Sigma_{X, 2k}$ are also well-separated. \square

The final two examples illustrate this proposition and provide explicit degree bounds on strict-separators on some smooth toric surfaces. Unlike for curves, our techniques do not typically prove that these degree bounds are sharp. However, for the classical case of ternary octics, we do obtain tight degree bounds for the existence of sum-of-square multipliers.

Example 5.16 (Strict-separators on toric surfaces of minimal degree). Let X be a toric surface of minimal degree. By combining Example 5.12 or Example 5.14 with Proposition 5.15, it follows that, for all $j \geq 2$, there exists elements $f \in P_{X,2j}$ such that, for all $k < j - 2$, the cones $\Sigma_{X,2j+2k}$ and $f \cdot \Sigma_{X,2k}$ are well-separated. In contrast, Example 4.17 only establishes that, for all $f \in P_{X,2j}$, the cones Σ_{X,j^2+j} and $f \cdot \Sigma_{X,j^2-j}$ are not well-separated, so there is a gap between our bounds. Since Example 4.17 also proves that, for all $f \in P_{X,2j}$, the cones $\Sigma_{X,4j-4}$ and $f \cdot P_{X,2(j-1)}$ are not well-separated, there is even a gap when we consider all nonnegative multipliers. \diamond

Proof of Theorem 1.2. Example 4.17 proves the first half and Example 5.16 proves the second. \square

Example 5.17 (Strict-separators on the projective plane). Let $Q := \text{conv}\{(0,0), (1,0), (0,1)\} \subset \mathbb{R}^2$ and let $\mathbb{P}^2 = Y_Q \subseteq \mathbb{P}^2$ be embedded by the very ample line bundle $A_Q = \mathcal{O}_{\mathbb{P}^2}(1)$. By combining Example 5.13 and Proposition 5.15, it follows that, for all $j \geq 2$, there exists elements $f \in P_{\mathbb{P}^2,2j}$ such that, for all $k < j - 3$, the cones $\Sigma_{\mathbb{P}^2,2j+2k}$ and $f \cdot \Sigma_{\mathbb{P}^2,2k}$ are well-separated. Example 4.18 shows that, for all $f \in P_{\mathbb{P}^2,8}$, the cones $\Sigma_{\mathbb{P}^2,12}$ and $f \cdot \Sigma_{\mathbb{P}^2,4}$ are not well-separated, so this degree bound for strict-separators on \mathbb{P}^2 is sharp when $j = 4$. \diamond

ACKNOWLEDGEMENTS

We thank Erwan Brugallé, Lionel Lang, and Mike Roth for useful conversations. The first author was partially supported by an Alfred P. Sloan Fellowship, the Simons Institute, and an NSF CAREER award DMS-1352073; the second author was partially supported by NSERC; and the third author was partially supported by the FAPA funds from Universidad de los Andes.

REFERENCES

- [AC+] Enrico Arbarello, Maurizio Cornalba, Phillip A. Griffiths, and Joe Harris, *Geometry of algebraic curves. Vol. I*, Grundlehren der Mathematischen Wissenschaften, vol. 267, Springer-Verlag, New York, 1985.
- [Be] Arnaud Beauville, *Complex algebraic surfaces*, 2nd ed., London Mathematical Society Student Texts, vol. 34, Cambridge University Press, Cambridge, 1996.
- [BI] Grigoriy Blekherman, *Positive Gorenstein ideals*, Proc. Amer. Math. Soc. **143** (2015), no. 1, 69–86.
- [BGP] Grigoriy Blekherman, João Gouveia, and James Pfeiffer, *Sums of Squares on the Hypercube*, to appear in Mathematische Zeitschrift, available at [arXiv:1402.4199](https://arxiv.org/abs/1402.4199) [math.AG].
- [BPT] Grigoriy Blekherman, Pablo A. Parrilo, and Rekha R. Thomas, *Semidefinite optimization and convex algebraic geometry*, MOS-SIAM Series on Optimization, vol. 13, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2013.
- [BSV] Grigoriy Blekherman, Gregory G. Smith, and Mauricio Velasco, *Sums of squares and varieties of minimal degree*, to appear in Journal of the American Mathematical Society, available at [arXiv:1308.0751](https://arxiv.org/abs/1308.0751) [math.AG].
- [Br] Erwan Brugallé, *Pseudoholomorphic simple Harnack curves*, to appear in L’Enseignement Mathématique, available at [arXiv:1410.2423](https://arxiv.org/abs/1410.2423) [math.AG].
- [C] David A. Cox, *The functor of a smooth toric variety*, Tohoku Math. J. (2) **47** (1995), no. 2, 251–262.
- [CLS] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [E] David Eisenbud, *The geometry of syzygies*, Graduate Texts in Mathematics, vol. 229, Springer-Verlag, New York, 2005.

- [EH] David Eisenbud and Joe Harris, *On varieties of minimal degree (a centennial account)*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), Proc. Sympos. Pure Math., vol. 46, Amer. Math. Soc., Providence, RI, 1987, pp. 3–13.
- [G] Gert-Martin Greuel, *SURFER 2012, ray-tracing software for visualizing real algebraic surfaces*. available at <https://imaginary.org/program/surfer>.
- [GLP] Laurent Gruson, Robert Lazarsfeld, and Christian Peskine, *On a theorem of Castelnuovo, and the equations defining space curves*, Invent. Math. **72** (1983), no. 3, 491–506.
- [Ha1] Joe Harris, *The genus of space curves*, Math. Ann. **249** (1980), no. 3, 191–204.
- [Ha2] ———, *A bound on the geometric genus of projective varieties*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **8** (1981), no. 1, 35–68.
- [Ha3] ———, *Curves in projective space*, Séminaire de Mathématiques Supérieures, vol. 85, Presses de l’Université de Montréal, Montréal, Que., 1982.
- [Hil] David Hilbert, *Über ternäre definite Formen*, Acta Math. **17** (1893), no. 1, 169–197.
- [IK] Anthony Iarrobino and Vassil Kanev, *Power sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics, vol. 1721, Springer-Verlag, Berlin, 1999.
- [J] Jean-Pierre Jouanolou, *Théorèmes de Bertini et applications*, Progress in Mathematics, vol. 42, Birkhäuser, Basel, 1983.
- [KO] Richard Kenyon and Andrei Okounkov, *Planar dimers and Harnack curves*, Duke Math. J. **131** (2006), no. 3, 499–524.
- [LPR] Henri Lombardi, Daniel Perrucci, and Marie-Françoise Roy, *An elementary recursive bound for effective Positivstellensatz and Hilbert 17-th problem*, available at [arXiv:1404.2338](https://arxiv.org/abs/1404.2338) [math.AG].
- [M] Juan C. Migliore, *Introduction to liaison theory and deficiency modules*, Progress in Mathematics, vol. 165, Birkhäuser Boston, Inc., Boston, MA, 1998.
- [N] U. Nagel, *Non-degenerate curves with maximal Hartshorne-Rao module*, Math. Z. **244** (2003), no. 4, 753–773.
- [P] Dmitrii V. Pasechnik, *An efficient sum of squares nonnegativity certificate for quaternary quartic*, available at [arXiv:1511.03473](https://arxiv.org/abs/1511.03473) [math.AG].
- [RG] Motakuri Ramana and A. J. Goldman, *Some geometric results in semidefinite programming*, J. Global Optim. **7** (1995), no. 1, 33–50.
- [R] Bruce Reznick, *Uniform denominators in Hilbert’s seventeenth problem*, Math. Z. **220** (1995), no. 1, 75–97.
- [Sc1] Claus Scheiderer, *Sums of squares on real algebraic curves*, Math. Z. **245** (2003), no. 4, 725–760.
- [Sc2] ———, *A Positivstellensatz for projective real varieties*, Manuscripta Math. **138** (2012), no. 1-2, 73–88.
- [Sc3] ———, *Semidefinite representation for convex hulls of real algebraic curves*, available at [arXiv:1208.3865](https://arxiv.org/abs/1208.3865) [math.AG].
- [ScR] Rolf Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, Encyclopedia of Mathematics and Its Applications, vol. 44, Cambridge University Press, Cambridge, 1993.

GREG BLEKHERMAN, SCHOOL OF MATHEMATICS, GEORGIA TECH, 686 CHERRY STREET, ATLANTA, GA, 30332, USA

E-mail address: greg@math.gatech.edu

GREGORY G. SMITH, DEPARTMENT OF MATHEMATICS AND STATISTICS, QUEEN’S UNIVERSITY, KINGSTON, ON, K7L 3N6, CANADA

E-mail address: ggsmith@mast.queensu.ca

MAURICIO VELASCO, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE LOS ANDES, CARRERA 1 NO. 18A 10, EDIFICIO H, PRIMER PISO, 111711 BOGOTÁ, COLOMBIA AND DEPARTAMENTO DE MATEMÁTICAS Y APLICACIONES, UNIVERSIDAD DE LA REPÚBLICA (CURE), MALDONADO, URUGUAY

E-mail address: mvelasco@uniandes.edu.co or mvelasco@cmat.edu.uy